

---

# Comparative Analysis of Ethanolic *Juniperus Thurifera* Leaf, Stem Bark and Root Extract Using Gas Chromatography and Mass Spectrometry

---

Singh Sharma<sup>1\*</sup>, Alagbe Olujimi John<sup>2</sup>, Liu Xing<sup>3</sup>, Sharma Ram<sup>4</sup>, Kumar Amita<sup>5</sup>

<sup>1\*,3,4,5</sup>Department of Microbiology, Sumitra Research Institute, Gujarat, India.

<sup>2</sup>Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Corresponding Email: <sup>1\*</sup>singh0081@gmail.com

**Received:** 02 August 2022    **Accepted:** 18 October 2022    **Published:** 22 November 2022

**Abstract:** Medicinal plants are reservoir of bioactive compounds or phytochemicals (terpenoids, alkaloids, tannins, flavonoids and so on) with marked pharmacological properties such as; antimicrobial, antioxidant, anti-inflammatory, antibacterial, analgesics, antiprotozoal, cytotoxic, anti-androgenic, antiviral, antipyretic, antitumor, anti-depressant, hypolipidemic and antihelminthic activities. This study was designed to examine the comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectrometry. Result obtained shows that *Juniperus thurifera* leaf extract contained 33 bioactive compounds which accounts to 75.47 % while *Juniperus thurifera* stem bark and root extract contained 37 and 50 phytochemicals with an aggregate of 80.77 % and 93.07 % respectively. Monoterpenes hydrocarbon (40.50 %), oxygenated monoterpenes (12.66 %), sesquiterpenes hydrocarbon (13.79 %) and oxygenated sesquiterpenes were found in *Juniperus thurifera* leaf extract while *Juniperus thurifera* stem bark and root extract contains monoterpenes hydrocarbon (48.72 %, 51.33 %), oxygenated monoterpenes (14.88 %, 20.49 %), sesquiterpenes hydrocarbon (10.71 %, 14.25 %) and oxygenated sesquiterpenes (6.56 %, 7.88 %) respectively. It was concluded that *Juniperus thurifera* leaf, stem bark and root ethanolic extract contains several secondary metabolites which can be used for the treatment of diseases without causing any deleterious effect on the environment and general performance of animals.

**Keywords:** Gas Chromatography, Mass Spectrometry, *Juniperus Thurifera*, Leaf, Root, Stem Bark, Phytochemicals.

## 1. INTRODUCTION

Medicinal plants are a rich source of phytochemicals and natural compounds (alkaloids, flavonoids, steroids, terpenoids, tannins and phenols) with marked therapeutic properties (Ruchika et al., 2022; Roma-Marzio et al., 2017). They are potential source of drugs and are

traditionally used in the treatment of digestive disorders, measles, cold, cough, diarrhea, arthritis, skin disease, joint pain, hemorrhoids, fever, sore throat and sexually transmitted infections (Misharina et al., 2009; Satrani et al., 2019). Medicinal plants are safe, environmentally friendly and performs antioxidant, antifungal, antiprotozoal, cytotoxic, hepato-protective, antibacterial, analgesics antipyretic, anti-inflammatory, antipyretic, antihelminthic, anti-fibrotic, hypolipidemic, anti-tumor and antioxidant activities (Cosentino et al., 1999; Alessandra et al., 2005).

There are over 200,000 species of medicinal plant with several pharmacological properties and many are still yet to be discovered (Alagbe, 2022). Unlike most synthetic or conventional drugs, medicinal plants are used to boost the immune system, suppress the activities of pathogenic organisms and maintain the general health status of animal without causing any deleterious effect (Olafadehan et al., 2021). In the midst of the prospective medicinal plant is *Juniperus thurifera* which is reported to contain several secondary metabolites necessary to bridge the gap between livestock production and food safety (Shittu et al., 2021).

*Juniperus thurifera* is an evergreen multipurpose small coniferous tree belonging to the family *capricidae* and class *pinopsida*. It is native to South Western Europe, North Africa and some parts of Asia including India (Farjon, 1992). The trees can grow up to 20 m tall with single or multiple trunks. The leaves are light green while the stems are grey brown at maturity with several bioactive compounds such as;  $\alpha$ -Murolene,  $\alpha$ -Longipinene,  $\gamma$ -eudesmol,  $\beta$ -Cayrophyllene,  $\alpha$ -Pinene, Torreyol- $\alpha$ -cadinol,  $\alpha$ -Cadinol,  $\alpha$ -Thujone,  $\beta$ -Phellandrene, Cis-thujanol and many more (Barrero et al., 2004). In vitro analysis has shown that ethanolic and methanolic extract from *Juniperus thurifera* leaf, stem bark and root can suppress the activities of some Gram +ve and -ve bacteria's such as; *Salmonella* spp, *Staphylococcus* spp, *Klebsiella* spp, *Micrococcus luteus*, *Escherichia coli*, *Citrobacter freundii*, *Bacillus subtilis*, *Neisseria* spp and *Enterobacter* spp thus preventing dysbiosis (Monsouri et al., 2010; Hajjouji et al., 2019). The use of gas chromatography and mass spectrometry will further help to identify some trace elements, gases as well as giving a clue on possible contaminants in the sample. Therefore, this experiment was designed to examine the comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectrometry.

### **Site of the experiment**

The experiment was carried out at the Department of Microbiology, Sumita Research Institute located within 23° 13'N 72°41'E Gujarat India in the month of September to November, 2022.

## **2. COLLECTION OF PLANT MATERIAL AND EXTRACTION METHODS**

Fresh *Juniperus thurifera* leaf, root and stem bark were harvested in Punsar village Gujarat. It was identified and authenticated by a Taxonomist (Dr. Singh Amit). It was washed separately with running tap water chopped into bits and allowed to dry under shade for 25 days. The dried samples were grinded separately using an electric blender. Dried *Juniperus thurifera* leaf, stem bark and root were soaked separately in ethanol (70 %) over a 48 hours period at a room temperature of 25°C with soxhlet apparatus. The mixture was stirred and filtered with Whatman filter paper, thereafter, it was dried in a rotary evaporator. The extract was stored individually a well labelled container and kept in the refrigerator at 4 °C before it was sent to the laboratory for additional examination. JTLE (*Juniperus thurifera* leaf extract), JTSBE (*Juniperus thurifera* stem bark extract) and JTRE (*Juniperus thurifera* root extract).

### Analysis of bioactive compounds in Juniperus thurifera leaf, stem bark and root extract

Secondary compounds in Juniperus thurifera leaf, stem bark and root extract were carried out using Labtron high precision Gas chromatograph mass spectrometer (Model GC-MS-879) with pre-filter mass analyzer and electron multiplier ensuring high sensitivity. Desorption of the extracts were performed at a temperature between 100 – 350 °C with a column flow rate of 1.0 -1.5 mL per minute with slit ratio of 1000: 1, pressure range (0-100 psi), temperature programming (7 stages/ 8 platforms) and electronic pressure control which supports CV and CC.

Table 1: Bioactive compounds in Juniperus thurifera leaf, stem bark and root extract using GC-MS

Bioactive compounds	% Area		
	JTLE	JTSBE	JTRE
$\gamma$ -Gurjunene	5.72	0.15	1.93
$\beta$ -Eudesmol	-	1.84	0.60
$\alpha$ -Cadinol	0.01	0.06	1.37
$\alpha$ -Humulene	1.12	1.51	2.05
$\delta$ -Cadinene	0.40	0.37	0.21
$\alpha$ -Cubebene	-	-	1.55
$\gamma$ -Cadinene	0.05	0.83	0.78
Germacrene D	-	0.20	1.49
Sabinene	15.88	17.03	20.82
$\gamma$ -Terpinene	0.01	0.88	1.53
$\alpha$ -Longipinene	4.02	5.06	1.07
$\alpha$ -Murolene	-	1.41	0.59
$\gamma$ -eudesmol	0.07	1.05	0.70
$\beta$ -Bourbonene	-	-	1.22
$\beta$ -Cayrophyllene	1.76	2.09	0.15
$\beta$ -Santalene	-	-	1.00
$\alpha$ -Pinene	16.81	19.97	24.31
$\beta$ -Citrylideneethanol	0.01	0.06	1.50
Torreyol- $\alpha$ -cadinol	0.03	0.01	0.07
D-Limonene	3.08	4.55	2.01
Elemol	-	-	0.01
Germacrene B	-	-	0.02
$\alpha$ -Cadinol	0.05	0.67	0.46
$\delta$ -2- Carene	0.93	0.55	1.01
$\delta$ -3- Carene	0.92	-	0.13
$\alpha$ -Pinene oxide	1.00	1.88	0.02
(Z)- $\beta$ - Ocimene	0.02	-	0.25
$\delta$ -Cadinene	0.57	1.21	1.40
$\beta$ -Eudesmol	0.61	0.40	1.22
Thymol	0.25	1.22	0.05
$\alpha$ -Phellandrene	1.06	0.02	0.09

$\alpha$ -Fenchene	0.10	0.08	0.17
Linalyl acetate	-	0.05	0.22
Citronellol	0.01	-	0.03
Terpinolene	1.88	2.33	2.41
Linalool	5.09	8.79	6.72
$\alpha$ -Thujone	2.00	1.72	2.60
$\alpha$ -Campholenal	1.36	0.58	0.07
Sabinene	-	0.27	0.44
Cis-sabinene hydrate	-	-	0.30
$\beta$ -Phellandrene	1.59	3.44	0.62
Cis-thujanol	4.02	5.07	0.88
Trans-4-thujanol	2.01	1.11	0.02
p-Cymen-8-ol	-	1.10	0.21
Trans-calamenene	3.03	-	0.45
Cis-murraola-(4)5-diene	-	-	1.82
Verbenone	-	-	0.52
$\alpha$ -Terpinyl acetate	-	2.03	0.01
Germacrene D-4-ol	-	0.01	2.03
Linalyl acetate	-	-	3.94
Aggregate (%)	75.47	80.77	93.07
<b>Breakdown (%)</b>			
Monoterpenes hydrocarbon (%)	40.50	48.62	51.33
Oxygenated monoterpenes (%)	12.66	14.88	20.49
Sesquiterpenes hydrocarbon (%)	13.79	10.71	14.25
Oxygenated sesquiterpenes (%)	8.83	6.56	7.88

JTLE: Juniperus thurifera leaf extract; TSBE: Juniperus thurifera stem bark extract and JTRE: Juniperus thurifera root extract




Figure 1: Graphical representation of bioactive compounds in Juniperus leaf, stem bark and root extract

### 3. RESULTS AND DISCUSSION

Table 1 revealed the gas chromatograph and mass spectrometry (GC-MS) analysis of bioactive compounds in *Juniperus thurifera* leaf, stem bark and root extract. Result obtained shows that *Juniperus thurifera* leaf extract contained 33 bioactive compounds (phytochemicals) which accounts to 75.47 % while *Juniperus thurifera* stem bark and root extract contained 37 and 50 phytochemicals with an aggregate of 80.77 % and 93.07 % respectively. Monoterpenes hydrocarbon (40.50 %), oxygenated monoterpenes (12.66 %), sesquiterpenes hydrocarbon (13.79 %) and oxygenated sesquiterpenes were found in *Juniperus thurifera* leaf extract while *Juniperus thurifera* stem bark and root extract contains monoterpenes hydrocarbon (48.72 %, 51.33 %), oxygenated monoterpenes (14.88 %, 20.49 %), sesquiterpenes hydrocarbon (10.71 %, 14.25 %) and oxygenated sesquiterpenes (6.56 %, 7.88 %) respectively. Secondary metabolites in the extracts was highest in *Juniperus thurifera* root extract, intermediate in *Juniperus thurifera* stem bark extract and lowest in *Juniperus thurifera* leaf extract. The result obtained is in agreement with the findings of Barrero et al. (2004); Alagbe et al. (2022) but contrary to the reports of Rachid et al. (2019). According to Oluwafemi et al. (2022); Agubosi et al. (2022) variations in the phytochemical components of medicinal plants could be attributed to species, age of plant, geographical location as well as the extraction procedure employed. Secondary metabolites or phytochemicals are chemicals of plant origin which possess pharmacological actions and are used by plants for defense against competitors, pathogens or predators (Alagbe and Ushie, 2022; Oluwafemi et al., 2022). They can also act as co-factors for enzymatic reactions and substrate for enzymes (Alagbe, 2022; Ramdani et al., 2013), ability to scavenge free radicals capable of causing diseases and selective inhibitors of

pathogenic organisms in the body (Hajjouji et al., 2019; Adewale et al., 2021). Terpenes are the largest group of bioactive compounds which may occur as oxygenated derivatives such as ketones, alcohols, aldehydes, phenol oxides and ketones (Muritala et al., 2022; Singh et al., 2021). Flavonoids are group of phenolic compounds with antioxidant, anti-allergic, antiviral, anti-tumor and anti-inflammatory activities (Shittu et al., 2021; Kumar and Pandey, 2013), prevent diseases caused by free radicals (Agubosi et al., 2022; Aoki et al., 2000) and prevents cardiovascular diseases (Oluwafemi et al., 2021; Takahashi and Ohnishi, 2004). Phenols are compounds possessing one or more aromatic rings with one or more hydroxyl groups. It has been reported to have antioxidant and anti-inflammatory activities (Olafadehan et al., 2021; Kim and Choi, 2013). Tannins are polyphenolic substance found in many plants product of secondary metabolism. They possess antibacterial, antiseptic and anti-carcinogenic properties (Lopez et al., 2008; Zheng and Wang, 2001), decrease in blood urea nitrogen and inhibition of lipid peroxidation (Amorati et al., 2013; Khan et al., 2009). Alkaloids are alkali-like substances with basic nitrogenous compounds of plant origin possessing a marked physiological function on animals (Hvattum, 2002; Giusti and Wrolstad, 2003).

#### 4. CONCLUSION

It was concluded that *Juniperus thurifera* leaf, stem bark and root extract contains several bioactive compounds which has marked therapeutic properties. It can also be used as natural alternative to antibiotics to bridge the gap between food safety and livestock production.

#### 5. REFERENCES

1. Muritala, Daniel Shittu., Alagbe, J.O., Ojebiyi, O.O., Ojediran, T.K and Rafiu, T.A. (2022). Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics. *Animal Science and Genetics* 18(2): 57-71.
2. Kumar S and Pandey, A.K. (2013) Chemistry and biological activities of flavonoids: an overview. *Scientific World Journal* 2013: 162750.
3. Aoki, T., Akashi, T and Ayabe, S. (2000) Flavonoids of leguminous plants: structure, biological activity, and biosynthesis. *Journal of Plant Research*, 113: 475–488.
4. Takahashi, A and Ohnishi, T. (2004). The significance of the study about the biological effects of solar ultraviolet radiation using the exposed facility on the international space station. *Biological Science Space* 18: 255–260.
5. Kim, S.H and Choi, K.C. (2013) Anti-cancer effect and underlying mechanism(s) of kaempferol, a phytoestrogen, on the regulation of apoptosis in diverse cancer cell models. *Toxicological Research*, 29: 229–234.
6. Zheng, W and Wang S.Y. (2001) Antioxidant activity and phenolic compounds in selected herbs. *Journal of Agriculture Food Chemistry*, 49: 5165–5170.
7. Khan, M.T., Orhan, I and Enol, S.S. (2009) Cholinesterase inhibitory activities of some flavonoid derivatives and chosen xanthone and their molecular docking studies. *Chemical Biological Interaction*, 181, 383–389.
8. Hvattum, E (2002) Determination of phenolic compounds in rose hip (*Rosa canina*) using liquid chromatography coupled to electrospray ionisation tandem mass spectrometry and diode-array detection. *Rapid Communication and Mass Spectrometer*, 16: 655–660.



9. Giusti, M and Wrolstad, R. (2003) Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal*, 14: 217–225.
10. Alagbe John Olujimi, Ramalan Sadiq Muhammad., Shittu Muritala Daniel and Olagoke Olayemi Christiana (2022). Effect of *Trichilia monadelpha* stem bark extract on the fatty acid composition of rabbit's thigh meat. *Journal of Environmental Issues and Climate Change* 1(1): 63-71.
11. Alagbe, J.O., Shittu, M.D and Tanimomo, Babatunde K. (2022). Influence of *Anogeissus leio carpus* stem bark on the fatty acid composition in meat of broiler chickens. *European Journal of Life Safety and Stability* 14(22): 13-22.
12. Alagbe, J.O (2022). Use of medicinal plants as a panacea to poultry production and food security: A review. *Gospodarka I Innowacje* 22(2022): 1-12.
13. Agubosi, O.C.P., Alexander, James and Alagbe, J.O. (2022). Influence of dietary inclusion of Sunflower (*Helianthus annus*) oil on growth performance and oxidative status of broiler chicks. *Central Asian Journal of Medical and Natural Sciences* 2(7): 187-195.
14. Agubosi, O.C.P., Soliu, M.B and Alagbe, J.O. (2022). Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. *Central Asian Journal of Theoretical and Applied Sciences* 3(3): 30-39.
15. Ruchika, G., Jelena, Z and Sekar, V. (2022). Medicinal plants and natural products in health promotion and disease prevention. *Evidence based Complementary and Alternative Medicine*, 22:283-291
16. Alagbe, J.O. (2022). Gas chromatography and mass spectroscopy of *Juniperus phoenicea* stem bark extract and its influence on the haemato-biochemical values of growing rabbits. *British Scientific Periodical* 1(1): 18-33.
17. Alagbe, J.O., Zubairu Habiba., Adedeji, O.M., Bamigboye, S and Dora Agbonika (2022). Influence of *Juniperus thurifera* root extract on the nutrient digestibility and caecal microbial count of growing rabbits. *Web of Synergy: International Interdisciplinary Research Journal* 1(1): 5-17.
18. Alagbe, J.O., Shittu, M.D., Ramalan, S.N., Tanimomo, K.B and Adekunle, D.A. (2022). Growth performance, semen quality characteristics and hormonal profile of male rabbit bucks fed *Rubia cordifolia* root extracts. *International Journal of Biological Engineering and Agriculture* 1(1): 1-13.
19. Agubosi, O.C.P., Imudia, Favour Dumkenechukwu and Alagbe, J.O. (2022). Evaluation of the nutritional value of air dried and sun-dried sweet potato (*Ipomoea batatas*) peels. *European Journal of Life Safety and Stability* 14(22): 43-51.
20. Alagbe, J.O., Agubosi, O.C.P., Oluwafemi, R.A and Gabriel Zakara (2022). Efficacy of *Trichilia monadelpha* stem bark on the growth performance of growing rabbits. *British Journal of Innovation in Science, Research and Development*, 1(2): 10-19.
21. Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Sciences* 3(2): 19-29.
22. Agubosi, O.C.P., Wika, B.K and Alagbe, J.O. (2022). Effect of dietary inclusion of Sunflower (*Helianthus annus*) oil on the growth performance of broiler finisher chickens. *European Journal of Modern Medicine and Practice*, 2(5): 1-10.



23. Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: performance and nutrient retention. *Discovery* 58(314): 134 -142.
24. Alagbe, J.O and Ushie, F.T. (2022). Growth performance of broiler chicks fed diets containing different levels of aqueous *Citrus aurantium* stem bark extracts. *Discovery* 58(319): 735-741.
25. Oluwafemi, R.A., Agubosi, O.C.P and Alagbe, J.O. (2021). Proximate, minerals, vitamins and amino acid composition of *Prosopis africana* (African mesquite) seed oil. *Asian Journal of Advances in Research* 11(1): 21-27.
26. Agubosi, O.C.P., Oluwafemi, R.A., and Alagbe, J.O. (2021). Preliminary study on GC-MS analysis of *Prosopis africana* seed (African mesquite) oil. *Journal of Ethics and Diversity in International Communication* 1(4): 18-20.
27. Alagbe, J.O., Adedeji, M.O., Habiba, Z., Nwosu, Gloria and Wyedia Dabara Comfort (2021). Physico-chemical properties of *Indigofera zollingeriana* seed oil. *Asian Journal of Advances in Medical Science* 3(4): 306-308.
28. Alagbe, J.O., Shittu, M.D and Ushie, F.T. (2021). GC-MS analysis of methanolic stem bark extract of *Zollingeriana indigofera*. *Asian Journal of Advances in Research* 11(4): 144-146.
29. Alagbe, J.O (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogetic Feed Additive in Growing Rabbit Diets: Growth Performance and Caecal Microbial Population. *Concept in Dairy and Veterinary Sciences*. 4(2):2021.
30. Adewale, A.O., Alagbe, J.O., Adeoye, Adekemi. O. (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogetic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 3(3): 1-12.
31. Singh, A.S., Alagbe, J.O., Sharma, S., Oluwafemi, R.A and Agubosi, O.C.P. (2021). Effect of dietary supplementation of melon (*Citrullus linatus*) seed oil on the growth performance and antioxidant status of growing rabbits. *Journal of Multidimensional Research and Reviews*, 2(1): 78-95.
32. Shittu, M.D., Alagbe, J.O., Adejumo, D.O., Ademola, S.G., Abiola, A.O., Samson, B.O and Ushie, F.T. (2021). Productive Performance, Caeca Microbial Population and Immune-Modulatory Activity of Broiler Chicks Fed Different Levels *Sida Acuta* Leaf Extract in Replacement of Antibiotics. *Bioinformatics and Proteomics Open Access Journal* 5(1): 000143.
33. Alagbe, J.O. (2021). *Prosopis africana* stem bark as an alternative to antibiotic feed additives in broiler chicks diets: Performance and Carcass characteristics. *Journal of Multidimensional Research and Reviews*, 2(1): 64-77.
34. Alagbe, J.O. (2021). *Daniellia oliveri* leaf extracts as an alternative to antibiotic feed additives in broiler chicken diets: Meat Quality and Fatty acid composition. *Indonesian Journal of Innovation and Applied Sciences* 1(3): 177-186.
35. Oluwafemi, R.A., Uankhoba, I.P and Alagbe, J.O. (2021). Effects of turmeric oil as a dietary supplements on the growth performance and carcass characteristics of broiler chicken. *International Journal of Orange Technologies*, 3(4): 1-9.





36. Oluwafemi, R.A., Uankhoba, I.P and Alagbe, J.O. (2021). Effects of turmeric oil as a dietary supplement on the haematology and serum biochemical indices of broiler chickens. *Bioinformatics and Proteomics Open Access Journal* 5(1): 000138.
37. Oluwafemi, R.A., Daniel, S.E and Alagbe, J.O. (2021). Haematology and serum biochemical indices of broiler chicks fed different inclusion levels of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil mixture. *International Journal of Discoveries and Innovations in Applied Sciences* 1(4): 20-26.
38. Oluwafemi, R.A., Abdullahi, H and Alagbe, J.O. (2021). Effect of dietary inclusion of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil mixture on the growth performance and caecal microbial population of broiler chickens. *Bioequivalence and Bioavailability International Journal* 5(2): 000153.
39. Alagbe, J.O (2020). Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and roots of *Indigofera tinctoria*. *International Journal on Integrated Education*. 3(10): 150-157.
40. Bonsignore, L., Loy, G., D. Secci, A. De Logu, G. A. (1990). Palmieri, Preliminary microbiological screening of Sardinian plants, *Fitoterapia*. 61: 339-341.
41. Zinoviadou, K.G., Koutsoumanis, K.P., Biliaderis, C.G. (2009). Physico-Chemical properties of whey protein isolate films containing oregano oil and their antimicrobial action against spoilage flora of fresh beef, *Meat Science*, 82: 338–345.
42. Cosentino, S., Tuberoso, C.I.G., Pisano, B., M. Satta, V. Mascia, E. Arzedi, F. Palmas. (1999). In vitro antimicrobial activity and chemical composition of Sardinian *Thymus* essential oils, *Letters of Applied Microbiology*, 29, 130-135
43. Alagbe, J.O., Ajagbe, A.D., Attama Jeremiah, Philemon, K.C and Bello, Kamoru, A (2020). *Albizia lebbek* stem bark aqueous extract as alternative to antibiotic feed additives in broiler chicks diets: Haematology, Serum indices and oxidative status. *International Journal of Biological, Physical and Chemical Studies*, 2(1): 8-15.
44. Alagbe, J.O (2020). Caecal Microbial Population of Growing Grass Cutters (*Thyronoymus Swinderianus*) Fed *Phyllanthus Amarus* and *Pilogstigma Thonngii* Leaf Meal Mixture as Partial Replacement for Soya Bean Meal. *Concept of Dairy and Veterinary Sciences*. 3(5): 350 – 355.
45. Alagbe, J.O., Zubairu Habiba., Adedeji, O.M., Bamigboye, S and Dora Agbonika (2022). Influence of *Juniperus thurifera* root extract on the nutrient digestibility and caecal microbial count of growing rabbits. *Web of Synergy: International Interdisciplinary Research Journal* 1(1): 5-17.
46. Olafadehan, O.A., Oluwafemi, R.A and Alagbe, J.O. (2020). Performance, haemato-biochemical parameters of broiler chicks administered Rolfe (*Daniellia oliveri*) leaf extract as an antibiotic alternative. *Advances in Research and Reviews*, 2020, 1:4.
47. Olafadehan, O.A., Oluwafemi, R.A and Alagbe, J.O. (2020). Carcass quality, nutrient retention and caeca microbial population of broiler chicks administered Rolfe (*Daniellia oliveri*) leaf extract as an antibiotic alternative. *Journal of Drug Discovery*. 14(33):146-154.
48. Alagbe, J.O. (2021). *Daniellia oliveri* leaf extracts as an alternative to antibiotic feed additives in broiler chicken diets: Meat Quality and Fatty acid composition. *Indonesian Journal of Innovation and Applied Sciences* 1(3): 177-186.
49. Rachid, Rahhal., Houda E.L Hajjouji., Said, Gmouh., Mohammed, Hsaine., Hassan, Fougrach and Wadi, Badri (2019). Chemical composition, antioxidant and antibacterial

- activities of the essential oils of *Juniperus phoenicea*, *Juniperus thurifera* and *Juniperus oxycedrus*. *Mediterranean Journal of Chemistry* 9(3): 190-198.
50. Farjon, A. (1992). The taxonomy of multiseed junipers (*Juniperus* Sect. *Sabina*) in southwest Asia and east Africa (Taxonomic notes on Cupressaceae I). *Journal of Botany*, 49: 251-283.
  51. Hammi, S., M., Ifriqui, V. Simonneaux L.Auclair. (2007). Évolution des recouvrements forestiers et de l'occupation des sols entre 1964 et 2002 dans la haute vallée des Ait Bouguemez (Haut Atlas Central, Maroc), *Sécheresse*. 18(4): 271–277.
  52. Barrero, A.F., Quilez, J.F., del Moral, M.M. Herrador, M. Akssira, A, Bennamara, S. Akkad., Aitigri, M. (2004). Oxygenated diterpenes and other constituents from Moroccan *Juniperus phoenicea* and *Juniperus thurifera* var. *Africana*, *Phytochemistry*, 65: 2507-2515.
  53. Adams, R.P. (1998). The leaf essential oils and chemotaxonomy of *Juniperus* sect. *Juniperus*, *Biochem Syst Ecology*, 26: 637-645.
  54. Misharina, T.A., Terenina, M.B and Krikunowa, N.I. (2009). Antioxidant properties of essential oils. *Applied Biochemistry and Microbiology*, 45: 642-647.
  55. El-Sawi, S.A., H.M. Motawae, A.M. Ali. (2007). Chemical composition, cytotoxic activity and antimicrobial activity of essential oils of leaves and berries of *Juniperus phoenicea* L. grown in Egypt, *African Journal of Traditional Complementary Alternative Medicine*, 4(4): 417-426.
  56. Ramdani, M., Lograda, T., H. Silini, A. Zeraib, P. Chalard, G. Figueredo, M. Bouchaala, S. Zerrar. (2013). Antibacterial Activity of Essential oils of *Juniperus Phoenicea* from Eastern Algeria, *Journal of Applied Pharmaceutical Science*. 3 (11): 022-028
  57. Hajjouji, H., C. Elagdi, R. Rahhal, S.Gmouh, M. Hsaine, H. Fougrach1, W. Badri. (2019). Chemical composition and antibacterial activity of essential oils extracted from three species of Moroccan *Juniperus*, *Research Journal of Pharmacy and Biological Sciences*, 10(3): 515-522.
  58. Hajdari, A., Mustafa, B., V. Gashi, D. Nebija, A. Ibraliu, J. Novak. (2014). Chemical composition of the essential oils of ripe berries of *Juniperus oxycedrus* L., growing wild in Kosovo, *Biochemical Systems and Ecology*, 57: 90-94.
  59. Satrani, B., Ghanmi, M., N. Mansouri, N. Amusant. (2015). Antioxidant properties of essential oils extracted from three species of Moroccan junipers, *Environmental Science Journal*, 11 (7): 239-247.
  60. Mansouri, N., Satrani, B., M. Ghanmi, L. El Ghadraoui, A. Aafi, A. Farah. (2010). Valorisation des huiles essentielles de *Juniperus thurifera* et de *Juniperus oxycedrus* du Maroc, *Phytothérapie*, 8: 166–170.
  61. Roma-Marzio, F., Najar, B., J. Alessandri, L. Pistelli and Peruzzi, L. (2017). Taxonomy of prickly juniper (*Juniperus oxycedrus* group): A phytochemical-morphometric combined approach at the contact zone of two cryptospecies, *Phytochemistry*, 141: 48-60.