

Research Paper



Studies on the effect of plant growth regulators on growth and yield of strawberry cv. (winter dawn) under protected cultivation

Ashish Ughareja¹, Yash Hemant Pandya^{2*}

^{1,2*}School of Agriculture, Lovely Professional University, Jalandhar, India.

Article Info

Article History:

Received: 03 December 2022

Revised: 12 February 2023

Accepted: 20 February 2023

Published: 08 April 2023

Keywords:

PGRs

Growth

Quality Fruit


Strawberry

Yield

Vegetative Growth

ABSTRACT

The experiment entitled “Studies on the Effect of Plant Growth Regulators on Growth and Yield of Strawberry cv. (Winter Dawn) Under Protected Cultivation” was conducted at experimental field of Department of Horticulture, Lovely Professional University, Phagwara, Punjab during 2021-22. The report’s findings showed that strawberry plants responded very well to the application of plant growth regulators. GA₃ 100 ppm gave the best result in vegetative growth, plant height, spread, leaves per plant, days taken to first flower and fruiting. When GA₃ 75 ppm gave best in fruit length, weight, diameter, Total soluble solid, total sugar, acidity of strawberry.



Corresponding Author:

Yash Hemant Pandya

School of Agriculture, Lovely Professional University, Jalandhar, India.

Email: wishyashpandya143@gmail.com

Copyright © 2023 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

The strawberry (*Fragaria x ananassa* Duch.) is an attractive Rosaceae fruit crop. This fruit is extremely valued in the international marketplace due to its appealing red colour, pleasant aroma, and excellent nutritional content in terms of vitamins (A and C) and minerals (Fe and K) [1]. Strawberry fruits,

in addition to being consumed fresh, are used to make jam and jelly because of the large content of pectin present [2]. *Fragaria x ananassa* Duch., the current cultivated strawberry, is a cross between two predominantly dioecious octoploid species, *Fragaria cheloensis* Duch and *Fragaria virginiana* Duch. Strawberry's chromosomal no is $2n=2x=56$. The species is octaploid. It's a short-lived (3-5 years), perennial, robust, stoloniferous herb that grows to a height of 10-20 cm and a spreading of 30-50 cm. The genuine strawberry fruit is an achene, a little, dry seed that is loosely linked to the swelling ovary wall. The strawberry's flesh is actually the ovary wall, which has many fruits/seeds on its surface.

Because the roots are shallow, plants require regular watering but not standing water [3]. Strawberry usually grows in temperate zones, and it is classified as a short-day plant based on its behaviour and eco system. Strawberry is a temperate plant that can be grown in both farmlands and mountains, but the fruit results are better on the hills. For best progress and growth, the strawberry needed a day temperature of 22°C and a night temperature of 7°C to 13°C. Frost and winter injuries decreased yield. Strawberry might cultivate in particular land, extending from thick clay to light sand and gravel. Strawberry plants, on the other hand, grow on sandy loam soil with a pH of 5.5 to 6.5. It is a heavy conveyor crop that produces more [4].

It is one of the few fruit trees that yields fast and easy and substantial returns per unit area on total income, as the fruit is ready for harvesting withing six months after sowing. Strawberry appears to be particularly responding to the implementation of plant growth regulators in addition to growth, production, and quality, according to research [1]. In India, strawberry fruit crop is still grown in open areas using a paddy straw mulching strategy by poor or marginal farmers, and it accounts for a large part of the national annual strawberry cultivable land. According to scientific findings, the strawberry plant responded positively to the application of a growth regulator [5]. Because of their suitability for treatment at a lower cost, naphthalene acetic acid (NAA) and gibberellins (GA_3) have been widely investigated in present agricultural systems. In many fruits, the role of these plant growth regulators has been explored [4]. Premature flowering, enhanced flowering time, collecting, and yield have all been observed with the use of GA_3 in strawberries. It boosts fruit output and quality, promotes cell elongation and expansion, helps to improve vegetative growth, reduces the period to maturity, and boosts fruit set [6]. In strawberry fruits, NAA promotes growth parameters, slows ripening, and enhances anthocyanin accumulation. It furthermore enhances the blooming period and improve fruit output and quality [6].

2. METHODOLOGY

The experiment was conducted during 2021-2022 at experimental filed of department of horticulture, Lovely Professional University, Phagwara, Punjab to experiment the influence of plant growth regulators on growth, yield, and quality of strawberry. The practical was carryout in a randomized block design with seven treatments consisted of control (no application of plant growth regulators), GA_3 (75, 100 ppm), NAA (75, 100 ppm), and CEPA (1000, 2000 ppm). All the treatment were replicated thrice. All the runners are equal and vigour transplanted during morning hours at a 30cm x 40cm. Daily watering was done for first week with drip irrigation and gap filling was done after first week of transplanted. After the transplant, the plant growth regulators were sprayed at 45, 60, and 75 days. In each replication, the observation was noted in three randomly selected plants. According to A.O.A.C [7], the quality of the fruits was tested during harvest using a standard method and procedure, and the results were statistically analysed.

2.1 Research Findings and Discussion

For this experiment, growth and flowering criteria including plant height, spread, number of leaves, number of flowers, number of days until first fruit, and the number of runners were chosen. The results of the current experiment make it evident that the application of the PGRs NAA, GA_3 , and CEPA had a major impact on the growth and flowering parameters. The maximum plant height (22.01 cm), plant spread (W-E) (24.16cm), plant spread (N-S) (23.49cm), no of leaves (22.97), days taken to first flower (41.04) no of flower (33.52), and days taken to first fruit (50.2) was recorded with treatment T_2 followed

by treatment T₁ whereas minimum result was observed in control T₀. When no of runners found best in treatment T₁. The number of fruits, flowers, and fruit productivity all significantly increased after GA₃ application, according to their observations of plant height, leaf count, leaf area, and fruit productivity. The evidence of [8], who also discovered that the application of GA₃ greatly improves plant height, leaf area, fruit setting percentage, and the number of runners is consistent with the study's findings as well.

Table 1. Effect of Plant Growth Regulators on Vegetative Growth and Flowering of Strawberry

Treatment	Plant Height (Cm)	Plant Spread (Cm) (W-E)	Plant Spread (Cm) (N-S)	No of Leaves	Days Taken to 1 st Flowering	No of Flower	Days Taken 1 st Fruit	No of Runners
T ₀ : Control	15.45	19.5825	20.275	18.4775	54.2	28.12	60.05	5.46
T ₁ : GA ₃ 75 ppm	21.125	22.425	23.09	20.8525	42.2	34.7	52.43	11.03
T ₂ : GA ₃ 100 ppm	22.01	24.1625	23.49	22.9775	41.04	33.52	50.2	9.12
T ₃ : NAA 75 ppm	18.38	21.5	22.28	20.6875	44.85	32.8	54.22	10.1
T ₄ : NAA 100 ppm	18.7375	21.9675	22.76	20.35	43.28	33.1	53.22	9.72
T ₅ : CEPA 1000 ppm	17.0475	20.43	20.525	20.015	46.03	31.25	55.12	7.21
T ₆ : CEPA 2000 ppm	17.4225	21.2075	20.5675	19.5675	47.3	30.2	56.38	6.31
C.D at 5%	1.79	2.078	2.098	1.97	4.407	3.070	5.256	0.824

2.2 Effect on Fruiting and Yield of Strawberry

The strawberry fruiting was adversely affected by the plant growth regulator treatments. The plant absorbs foliar spray of 75 ppm GA₃ produce maximum no of fruit (28.1), fruit length (4.81 cm), fruit weight (14.35 gm), fruit diameter (4.01 cm), fruit set (83.86 %), and yield were recorded in treatment T₁ followed by treatment T₂. When minimum observation was found in control T₀. In foliar spray with GA₃, additional biomass may be able to generate extra metabolites during photosynthesis, which eventually sank into the producing fruits and generated berry with the most weight. In strawberries, the use of GA₃ has been observed to boost berry weight [5]. A higher number of marketable fruits were produced as a result of the exogenous application of GA₃, which also indirectly affected the auxin metabolism and increased the fruit yield. It has also been previously documented that applying GA₃ increases fruit yield in strawberry [3], [9].

Table 2. Effect of Plant Growth Regulators on Yield Attributing and Yield of Strawberry

Treatment	No of Fruits	Fruit Length (Cm)	Fruit Weight (Gm)	Fruit Diameter (Cm)	Fruit Set (%)	Yield Per Plant (Gm)
T ₀ : Control	19.02	2.83	10.32	2.12	67.63869	196.286
T ₁ : GA ₃ 75 ppm	28.1	4.81	14.35	4.01	83.86167	417.585
T ₂ : GA ₃ 100 ppm	26.82	4.01	13.88	3.45	80.01193	372.262
T ₃ : NAA 75 ppm	25.43	3.8	13.26	2.98	77.53049	337.202
T ₄ : NAA 100 ppm	26.2	3.26	12.96	2.76	79.15408	339.552
T ₅ : CEPA 1000 ppm	23.18	3.1	11.38	2.43	74.176	263.788

T ₆ : CEPA 2000 ppm	22.11	3.01	10.81	2.21	73.21192	245.928
C.D at 5%	2.349	0.343	1.198	0.279	7.344	30.224

2.3 Effect on Fruit Quality of Strawberry

In terms of strawberry quality, the results reported in Table 3 demonstrated that the use of growth regulators altered the quality features of the strawberry fruit. The plant absorbs foliar spray of 75 ppm GA₃ produce maximum Total soluble solid (8.44 °B), Acidity (0.52%), Ascorbic acid (65.47mg), Total sugar (9.48 %), Reducing sugar (5.1%), non-reducing sugar (4.38%) were recorded in treatment T₁. When minimum result show in control T₀. Gibberellic acid was applied to fruits, which massively improved total soluble solids and decreased titratable acidity. With foliar treatment of 75 ppm GA₃, the highest total soluble solid was recorded. However, in the current investigation, the control group had the lowest levels of total soluble solids [10]. These results support those of [11], who demonstrated that GA₃ concentrations responded favourably to a strawberry quality measure.

Table 3. Effect of Plant Growth Regulators on Quality of Strawberry

Treatment	Total Soluble Solid (°B)	Acidity (%)	Ascorbic Acid (Mg/100 G)	Total Sugar (%)	Reducing Sugar (%)	Non-Reducing Sugar (%)
T ₀ : Control	6.9	0.79	52.69	6.03	3.9	2.13
T ₁ : GA ₃ 75 ppm	8.44	0.52	65.47	9.48	5.1	4.38
T ₂ : GA ₃ 100 ppm	7.91	0.54	63.11	8.35	4.31	4.04
T ₃ : NAA 75 ppm	7.49	0.71	59.84	8.1	4.12	3.98
T ₄ : NAA 100 ppm	8.01	0.68	57.14	8.89	4.72	4.17
T ₅ : CEPA 1000 ppm	7.32	0.76	55.54	7.8	4.03	3.77
T ₆ : CEPA 2000 ppm	7.1	0.71	54.22	7.2	3.98	3.22
C.D at 5%	0.730	0.070	5.609	0.764	0.417	0.356

3. CONCLUSION

Based on the data reviewed for this study, it is concluded that applying 75 ppm and 100 ppm GA₃ before flowering improved strawberry cv. (Winter dawn). overall vegetative development, flowering, fruit set, fruit size, yield, and fruit quality. These results could be utilised to assess the impact of PGR on various strawberry cultivars grown under protected agriculture in the future.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author Contributions Statement

Name of Author	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Ashish Ughareja	✓	✓	✓	✓		✓		✓	✓	✓	✓		✓	
Yash Hemant Pandya		✓		✓	✓		✓	✓		✓		✓	✓	✓

C : Conceptualization	I : Investigation	Vi : Visualization
M : Methodology	R : Resources	Su : Supervision
So : Software	D : Data Curation	P : Project administration
Va : Validation	O : Writing - Original Draft	Fu : Funding acquisition
Fo : Formal analysis	E : Writing - Review & Editing	

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study and their voluntary consent was obtained prior to data collection.

Ethical Approval

Not Applicable.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.



REFERENCES

- [1] Kumar, R., Bakshi, M., & Singh, D. B. (2012). Influence of plant growth regulators on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch.) under UP sub tropics. *Asian J. Hortic*, 7, 434-436.
- [2] K. D. Rathod, T. R. Ahlawat, S. Kumar, M. Sarkar, and B. Chakraborty, 'Effect of Plant Growth Regulators on Growth, Yield and Quality of Strawberry (*Fragaria x ananassa* Duch.) cv. Winter Dawn under Open Field Conditions of South Gujarat', *Agricultural Science Digest*, vol. 41, no. 2, pp. 329-333, 2021. doi.org/10.18805/ag.D-5240
- [3] Vishal VC, Thippesha D, Chethana K, Maheshgowda BM, Veerasha BG, Basavraj AK. Effect of various growth regulators on vegetative parameters of strawberry (*Fragaria x ananassa* Dutch) cv. Sujatha. *Research Journal of Chemical and Environmental Sciences*. 2016; 4:68-71.
- [4] Yadav, A., Pratap, B., Shivam, A. K., & Patro, A. (2018). Assess the effect of micronutrients and plant growth regulators on quality parameters of strawberry cv. Chandler. *The Pharma Innovation J*, 7(1), 303-305.
- [5] Mir MM, Barche S and Singh DB, Effect of plant growth regulators on growth, yield and quality of strawberry (*Fragaria x ananassa* Duch.) cv. 'Sweet Charlie'. *Applied Biological Research*, 2004, 6(1/2), p 48-51.
- [6] R. R. Sharma and R. Singh, 'Gibberellic acid influences the production of malformed and button berries, fruit yield and quality in strawberry (*Fragaria x ananassa* Duch)', *Scientia Horticulture*, vol. 119, no. 4, pp. 430-433, 2009. doi.org/10.1016/j.scienta.2008.11.002
- [7] A.O.A.C (1990). Official method analysis, Association of Analytical Chemist. Washington, D.C., U.S.A.
- [8] Qureshi, K. M., S. Chughtai, U. S. Qureshi and N. A. Abbasi (2013). Impact of exogenous application of salt and growth regulators on growth and yield of strawberry. *Pakistan Journal of Botany*, 45(4): 1179-1185.
- [9] R. R. Sharma and R. Singh, 'Gibberellic acid influences the production of malformed and button berries and fruit yield and quality in strawberry (*Fragaria x ananassa* Duch)', *Scientia Horticulturae*, vol. 119, pp. 430-433, 2009. doi.org/10.1016/j.scienta.2008.11.002
- [10] K. D. Rathod, T. R. Ahlawat, S. Kumar, M. Sarkar, and B. Chakraborty, 'Effect of Plant Growth Regulators on Growth, Yield and Quality of Strawberry (*Fragaria x ananassa* Duch.) Cv. Winter Dawn under Open Field Conditions of South Gujarat', *Agricultural Science Digest*, vol. 41, no. 2, pp. 329-333, 2021. doi.org/10.18805/ag.D-5240

- [11] Prasad, M., M. Minz, K. K. Jha, R. Kumar and B. Das (2013). Studies on the effect of mulching and pgrs on physicochemical characters and postharvest performance of strawberry (*Fragaria x ananassa* Duch.) cv. Douglas. *Journal of Interacademia*, 17(1) : 11-16.

How to Cite: Ashish Ughareja, Yash Hemant Pandya (2023). Studies on the effect of plant growth regulators on growth and yield of strawberry cv. (winter dawn) under protected cultivation, *International Journal of Agriculture and Animal Production (IJAAP)*, 3(1), 69-74. <https://doi.org/10.55529/ijaap.32.33.38>

BIOGRAPHIES OF AUTHORS

	<p>Ashish Ughareja, is a researcher, from the School of Agriculture at Lovely Professional University Jalandhar, India. His interest's kind of revolve around agronomy, sustainable farming systems, soil health management, and crop improvement practices. Lately he is actively involved in academic and research activities that aim to boost agricultural productivity using innovative and eco-friendly approaches, even if it sounds simple.</p>
	<p>Yash Hemant Pandya, is also linked with the School of Agriculture at Lovely Professional University, Jalandhar, India. His research interests include precision agriculture, plant science, and sustainable crop production. He takes part in research and academic initiatives that work on creating more efficient farming practices, and at the same time supporting sustainable rural development.</p>