

Research Paper



Investigation of dentinal tubule occlusion by a desensitizing dentifrice: an in-vitro SEM analysis

Lynn Johnson^{1*}, Sonal GP Pillai², Palak Upadhyay³

¹Associate Professor, Rama Dental College, Kanpur, UP, India.

²Ex-student, Rungta College of Dental Science and Research Centre, Bhilai, CG, India.

³Associate Professor, Maitri College of Dentistry & Research Centre, Durg, CG, India.

Article Info

Article History:

Received: 02 March 2023

Revised: 19 May 2023

Accepted: 25 May 2023

Published: 12 July 2023

Keywords:


Desensitizing Agents

Dentin Tubules

Desensitizing Agents

Dentine Sensitivity

Pro-Argin Technology



ABSTRACT

Background: Exposed and open dentinal tubules cause dentin hypersensitivity, which is a common clinical condition, and agents which can occlude dentinal tubules are of great therapeutic interest in this condition.

Objective: The aim of this study was to compare the dentin tubule-occluding effect of a dentifrice containing Pro-Argin Technology (Colgate Sensitive Plus) to patent dentin tubule-occluding effect of Pro-Argin Technology.

Methods: The methods involved preparation of twenty dentin discs and random assignment of these to two groups: one consisting of untreated (control) discs and the other consisting of dentin discs treated with the Pro-Argin dentifrice. The diameter of dentin tubules was examined and measured in both groups by scanning electron microscopy (SEM).

Results: Clear morphological differences between the two groups were determined by SEM imaging. Dentin tubules were mostly open and patent in the untreated control discs. On the contrary, all the dentifrice treated samples showed the presence of material on the dentin surface, filling the dentinal tubules, and to a large extent reducing and/or closing the dentin tubule openings.

Conclusion: From the results of the SEM analysis and relevant statistical analysis, it is concluded that the dentifrice containing Pro-Argin Technology has an ability to occlude dentinal tubules in vitro. The results provide further evidence of the effectiveness of Pro-Argin Technology as a dentin tubule occlusion product and demonstrate its potential as an anti-sensitivity agent in dental care products designed to manage dentin hypersensitivity.

Corresponding Author:

Lynn Johnson

Associate Professor, Rama Dental College, Kanpur, UP, India.

Email: lynnjohnson380@gmail.com

Copyright © 2023 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

The term "dentinal hypersensitivity" (DH) refers to a type of discomfort that results from exposed dentin and is generally brought on by chemical, thermal, tactile, or osmotic stimuli and cannot be attributed to any other type of dental disease or defect. [1], [2] Patients' subjective experiences with DH pain might differ greatly from one patient to the next. 33% to 88% of people have DH, which can cause varied degrees of discomfort when eating, drinking, or even breathing, according to Splieth and Tachou, [3], [4], [5] Having exposed dentine surfaces, open tubule orifices on the exposed dentine surface, and patent tubules leading to vital pulp are the three key diagnostic indicators of DH. Dentine exposure frequently results from the erosion of the cervical cementum, gingival recession, and erosion of the enamel brought on by various forms of tooth wear. [6], [7], [8] the commonly recognized view of the mechanism underlying DH states that following stimulation, fluid is displaced inside the dentinal tubules, causing pain.

Pain is caused by the activation of the nerve endings at the pulp-dentine interface caused by the movement of fluid inside and outside the tubules. [9], [10] The cause of dentine hypersensitivity and the characteristics of the lesion are currently only vaguely understood. The activation of pulpal mechanoreceptors is attributed to fluid movement inside exposed dentinal tubules, in accordance with the hydrodynamic hypothesis. Attrition resulting from occlusal discord, gingival recession following either a periodontal disease process or periodontal treatment, and trauma from tooth brushing are the most common causes of dentine exposure. For the treatment of hypersensitive dentine, a variety of chemicals have been explored thus far with varied degrees of effectiveness. The most popular treatments for reducing hypersensitivity include the use of iontophoresis, hard and soft tissue lasers, and dentifrices and mouthwashes containing desensitizing chemicals. [11] Unfortunately, the methods used to cure this issue are ineffective and have no long-term consequences. [12] This in-vitro SEM study's objective is to assess how well dentifrices containing pro-argin technology obstruct dentinal tubules to lessen dentinal hypersensitivity.

Resources and Procedures 20 maxillary first premolars that had just had orthodontic treatment were removed, cleaned, and kept in distilled water pending usage. The cementum layer was completely removed by scaling and root planing, revealing the dentine below. The cervical portion of the root was then prepped for a 2x2x1mm dentine disk utilizing a double diamond disk. The samples were then washed with distilled water after being treated with 17% ethylene-diamine-tetraacetic acid (EDTA) (Endo L, MAARC, Thane, Maharashtra, India) for 1 min to remove the smear layer. The samples were then split into two groups, each with ten samples: Control group in Group I; pro-argin group in Group II.

The diameter of untreated open dentine tubules in the control group served as the reference. The research group only received one application of the dentifrice, which was left on for two minutes and then washed off with distilled water. The samples were then fixed in ethanol solutions of increasing strength, dried, and gold-sputter coated in accordance with SEM specifications. Each specimen surface that was investigated using a SEM at magnifications of 500x, 1000x, and 2000x yielded photomicrographs. One researcher conducted a single blind analysis of the samples at a magnification of 2000x and determined the decrease in tubule diameter between the study group and the control group. The data for the control group's tubule diameter and the study group's tubule diameter decrease (measured in microns) were recorded on a data sheet. An unpaired t-test was then used to compare the two groups side by side, and the results were tallied.

1.1 Observation and Results

The mean tubule diameter for the control group was $1.97 \pm 0.49 \mu\text{m}$. Mean tubule diameter reduction achieved in the tooth-paste group was $0.83 \pm 0.23 \mu\text{m}$ ($42.1 \pm 11.8\%$). The unpaired t-test revealed extreme statistical difference in mean between the test and the control group ($p < 0.0001$). Thus,

the test sample showed significant reduction in the mean tubule diameter as compared to the control group Table 1.




Figure 1. Showing Open Dentinal Tubules in Control Group Under SEM 2000x Magnification




Figure 2. SEM Images of Partially and Completely Occluded Dentinal Tubules in the Test Group At 2000x Magnification

Table 1. Showing Dentinal Tubule Diameter Reduction of the Test Group with Reference to the Control Group

Mean	Control Group (C)	Tooth-Paste Group (TP)	
	Diameter Of Open Dentinal Tubule	Reduction Obtained (In Microns)	Reduction Obtained (In%)
	1.97	0.83	42.1
Standard Deviation (SD)	0.49	0.23	11.8

2. RESULTS AND DISCUSSION

It is accurate to infer that people will have their natural dentition in the oral cavity for a longer length of time as life expectancy rises and thus DH will likely become more common. Concomitantly, new eating practices, such as regularly consuming acidic foods and drinks, can cause an increase in tooth wear and, as a result, dentin hypersensitivity. [13] The body of research demonstrates that DH significantly

lowers patients' quality of life. [14] According to the Brännström and Aström hydrodynamic hypothesis, open dentinal tubules allow fluid to pass through them, stimulating the tooth pulp's nerve terminals. Scanning electron microscopy has revealed that tubules in clinically identified, "sensitive," exfoliated teeth are eight times more numerous, two times broader in diameter, and more permeable, in contrast to tubules in "non sensitive" teeth, which are less frequent, smaller, and typically plugged, as demonstrated in the research. [15] Dentin hypersensitivity decreases when dentinal fluid flow declines, in accordance with the rules of hydrodynamics. [16] Desensitizers with neural or obliterating effect are two options for treating DH. Products having neurological action, such potassium nitrate, oxalates, and low-power laser treatment (photobiomodulation), aim to suppress neural activity, which causes depolarization of nerve fibers.

Tubular occlusion can be treated with a variety of methods, including glutaraldehyde, varnishes, oxalates, biogas, and high-power lasers. These therapeutic approaches can also be categorized as at-home, over-the-counter desensitizing mouthwashes, dentifrices, and professional treatments. Patients favor home use products the most because they are the most practical, reasonable, and routine solutions for mild to moderate DH. These are also acceptable due to their accessibility and simplicity of use when combined with regular oral hygiene practices like cleaning teeth.2 Frequently used desensitizing substances can also be divided into four groups: tubule-occluding substances (calcium hydroxide, potassium nitrate, sodium fluoride), protein precipitants (formaldehyde, silver nitrate, and strontium chloride hexahydrate), anti-inflammatory substances (corticosteroids), and resins and adhesives. None of these substances can, however, have long-lasting effects since DTs would eventually be exposed again due to DTs' abrasion and erosion by internal and external acids. [17] Desensitizing mouthwashes and toothpastes work by precipitating crystals from their active components, which reduce dentinal tubule width.2 to treat this disease, there are several desensitizing toothpastes on the market right now. The majority of these products either block the exposed dentine tubules or desensitize the pulp nerve in order to function.

[18] With its unique pro-argin technology, the dentifrice Colgate Sensitive Plus utilized in this study is claimed to provide fast pain relief by constructing a calcium-rich coating that plugs the exposed dentine channels to shield sensitive teeth from unexpected shocks of pain. The open tubules are physically blocked and closed by the combination of arginine and calcium carbonate, according to early research, which results in desensitization of DH. Our findings also provide credence to the idea that arginine-rich toothpaste might reduce DH symptoms by shrinking the tubule diameter. [18] The findings of this investigation are consistent with those of other studies using scanning electron microscopy that were conducted on tooth paste containing pro-argin.

3. CONCLUSION

As a result of either lowering or blocking the exposed dentin tubules, the test dentifrice appears to be a useful tool for successfully treating dentine hypersensitivity, as shown by the results of the current investigation. To assess the long-term stability of the acquired favorable outcomes, more research is required, particularly long-term in-vitro and in-vivo investigations.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author Contributions Statement

Name of Author	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Lynn Johnson	✓	✓	✓	✓		✓		✓	✓	✓	✓	✓		✓
Sonal GP Pillai	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Palak Upadhyay	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

C : Conceptualization	I : Investigation	Vi : Visualization
M : Methodology	R : Resources	Su : Supervision
So : Software	D : Data Curation	P : Project administration
Va : Validation	O : Writing - Original Draft	Fu : Funding acquisition
Fo : Formal analysis	E : Writing - Review & Editing	

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study and their voluntary consent was obtained prior to data collection.

Ethical Approval

Not Applicable.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.


REFERENCES


- [1] H. Zhuang, Y. Liang, S. Xiang, H. Li, X. Dai, and W. Zhao, 'Dentinal tubule occlusion using Er:YAG Laser: an in vitro study', J. Appl. Oral Sci., vol. 29, p. e20200266, Apr. 2021. doi.org/10.1590/1678-7757-2020-0266
- [2] A. C. C. Aranha and C. de P. Eduardo, 'Effects of Er:YAG and Er,Cr:YSGG lasers on dentine hypersensitivity. Short-term clinical evaluation', Lasers Med. Sci., vol. 27, no. 4, pp. 813-818, July 2012. doi.org/10.1007/s10103-011-0988-9
- [3] R. Birang, J. Poursamimi, N. Gutknecht, F. Lampert, and M. Mir, 'Comparative evaluation of the effects of Nd:YAG and Er:YAG laser in dentin hypersensitivity treatment', Lasers Med. Sci., vol. 22, no. 1, pp. 21-24, Mar. 2007. doi.org/10.1007/s10103-006-0412-z
- [4] V. Maximiano, A. C. Machado, M. L. Yoshida, C. M. Pannuti, T. Scaramucci, and A. C. C. Aranha, 'Nd:YAG laser and calcium sodium phosphosilicate prophylaxis paste in the treatment of dentin hypersensitivity: a double-blind randomized clinical study', Clin. Oral Investig., vol. 23, no. 8, pp. 3331-3338, Aug. 2019. doi.org/10.1007/s00784-018-2759-5
- [5] M. B. Blatz, 'Laser therapy may be better than topical desensitizing agents for treating dentin hypersensitivity', J. Evid. Based. Dent. Pract., vol. 12, no. 3 Suppl, pp. 229-230, Sept. 2012. [doi.org/10.1016/S1532-3382\(12\)70044-1](https://doi.org/10.1016/S1532-3382(12)70044-1)
- [6] M.-L. Hu, G. Zheng, Y.-D. Zhang, X. Yan, X.-C. Li, and H. Lin, 'Effect of desensitizing toothpastes on dentine hypersensitivity: A systematic review and meta-analysis', J. Dent., vol. 75, pp. 12-21, Aug. 2018. doi.org/10.1016/j.jdent.2018.05.012
- [7] G. R. Holland et al., 'A short term comparative evaluation of the efficacy of diode laser with desensitizing toothpastes and mouthwashes in the treatment of dentinal hypersensitivity', J Clin Periodontol, vol. 24, no. 3, pp. e229-234, 1997.
- [8] K. Khare, P. Bhusari, A. Soni, S. K. Malagi, D. Abraham, and L. Johnson, 'A comparative evaluation of efficacy of BioMin F and propolis containing toothpastes on dentinal tubule occlusion with and without use of an adjunct 810 nm diode laser: An In vitro scanning electron microscope study', J Microsc Ultrastruct, vol. 11, pp. 41-46, 2023. doi.org/10.4103/jmau.jmau.133.20

- [9] D. H. Pashley, L. Rimondini, C. Baroni, and A. Carrassi, 'Ultrastructure of hypersensitive and non-sensitive dentine. A study on replica models', *J Clin Periodontol*, vol. 39, pp. 899-902, 1994. doi.org/10.1111/j.1600-051X.1995.tb01792.x
- [10] E. G. Absi, M. Addy, and D. Adams, 'Dentine hypersensitivity. A study of the patency of dentinal tubules in sensitive and nonsensitive cervical dentine', *J Clin Periodontol*, vol. 14, pp. 280-284, 1987. doi.org/10.1111/j.1600-051X.1987.tb01533.x
- [11] P. M. Bartold, 'Dentinal hypersensitivity: a review', *Aust Dent J*, vol. 51, no. 3, pp. 212-218, Sept. 2006. doi.org/10.1111/j.1834-7819.2006.tb00431.x
- [12] M. Brännström and A. Aström, 'The hydrodynamics of the dentine; its possible relationship to dentinal pain', *Int Dent J*, vol. 22, pp. 219-227, 1972.
- [13] D. H. Pashley, 'Mechanisms of dentin sensitivity', *Dent Clin North Am*, vol. 34, pp. 449-473, 1990. [doi.org/10.1016/S0011-8532\(22\)01128-4](https://doi.org/10.1016/S0011-8532(22)01128-4)
- [14] J. S. Earl, M. B. Ward, and R. M. Langford, 'Investigation of dentinal tubule occlusion using FIB-SEM milling and EDX', *J Clin Dent*, vol. 21, no. 2, pp. 37-41, 2010.
- [15] I. Petrou et al., 'A breakthrough therapy for dentin hypersensitivity: how dental products containing 8% arginine and calcium carbonate work to deliver effective relief of sensitive teeth', *J Clin Dent*, vol. 20, no. 1, pp. 23-31, 2009.
- [16] D. Cummins, 'Recent advances in dentin hypersensitivity: clinically proven treatments for instant and lasting sensitivity relief', *Am J Dent*, vol. 23, May 2010.
- [17] Cummins D. Recent advances in dentin hypersensitivity: clinically proven treatments for instant and lasting sensitivity relief. *Am J Dent*. 2010 May;23 Spec No A:3A-13A. PMID: 21284246.
- [18] S. A. Lavender et al., 'Mode of action studies on a new desensitizing dentifrice containing 8.0% arginine, a high cleaning calcium carbonate system and 1450 ppm fluoride', *Am J Dent*, vol. 23, May 2010.

How to Cite: Lynn Johnson, Sonal GP Pillai, Palak Upadhyay. (2023). Investigation of dentinal tubule occlusion by a desensitizing dentifrice: an in-vitro SEM analysis. *Journal of Nursing Research, Patient Safety and Practice (JNRPS)*, 3(2), 11-17. <https://doi.org/10.55529/jnrps.34.11.16>

BIOGRAPHIES OF AUTHORS

	Lynn Johnson, I believe an Associate Professor at Rama Dental College, Kanpur, Uttar Pradesh, India. Her academic and professional focus seems to sit mostly on clinical dentistry and dental research, with a strong pull towards prevention and restorative approaches for oral health. She has taken part in research related to dentistry and she is also actively involved in teaching, plus the hands-on clinical training of dental students. Overall her work shows a clear curiosity about evidence-based methods for better dental care, and for improving patient outcomes. Email: lynnjohnson380@gmail.com
	Sonal GP Pillai, is a dental professional and also a former student of Rungta College of Dental Science and Research Centre, Bhilai, Chhattisgarh, India. Her academic interests revolve around clinical dentistry and oral health research. She has joined in dental research work that involves evaluating and managing everyday oral and dental problems. In short her academic pathway reflects a wish to build more evidence-informed strategies for dental treatment, and to better patient support. Email: gsonalpillai@gmail.com



Palak Upadhyay, is an Associate Professor at Maitri College of Dentistry & Research Centre, Durg, Chhattisgarh, India. Her academic and professional interests include clinical dentistry, dental education, and oral health research. She contributes through teaching and other academic activities, and she also works on research aimed at refining diagnostic and therapeutic approaches within dentistry. Her overall direction, honestly, puts weight on evidence-based dental practice and on moving oral healthcare forward. Email: drpalak22@gmail.com