

Research Paper



Cervical cancer to see the evaluation of the effectiveness of colposcopy in iraqi women

Dr. Muna Kadhum Farhood Malla^{1*}, Dr. Massar Raheem Abbas², Dr. Zena Hataf Salih³

¹M. B. Ch. B \ D. G. O. \ (Obstetrics and Gynecology) Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq.

²M. B. Ch. B \ D. G. O. \ (Obstetrics and Gynecology) Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq.

³M. B. Ch. B. \ D. G. O. \ (Obstetrics and Gynecology) Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq.

Article Info

Article History:

Received: 11 January 2023

Revised: 30 March 2023

Accepted: 07 April 2023

Published: 25 May 2023

Keywords:


Cervical Cancer

Cervical Biopsy Swede

Scores and CIN

Colposcopy

Swede Score



ABSTRACT

Background: One of the leading causes of female death in developing nations is cervical cancer, which accounts for one-fourth of all cases globally. **Objective:** This paper was focused to assess the outcomes of cervical cancer to see the evaluation of the effectiveness of colposcopy in Iraqi women. **Patients and Methods:** This paper was represented as a cross-sectional study to assess the outcomes of cervical cancer to see the evaluation of the effectiveness of colposcopy in Iraqi women where to include from 25th August 2021 to 16th July 2022 into, 50 patients for ages between 25-50 years. This paper was analysed and examine of clinical features for cervical cancer by the SPSS program. **Discussion:** The specificity of the Swede score was scoring with seven or above was 87% in the current investigation, while the sensitivity for high-grade lesions was 92%. The threshold was lowered to 5, which boosted the sensitivity reached 72.42% and the specificity to 82%. Using a threshold of 5 for the Swedish score, the accuracy of the value of positive prediction is 75.48%. The accuracy considerably increased to 93.28% if the cutoff for the Swede score was set at 7. The results were similar to those that Bowring et al., as well as Strander et al., had previously reported. **Conclusion:** This study shows the Swede score was good as screening modalities as performance assessed of patients who have lower abdominal pain and discharge per vaginam as well as dyspareunia, abnormal uterine bleeding which colposcopy conducted into pointing of suspicious cervix with the unhealthy case. In comparison between the Swede score >5 and >7, our study noticed Swede score have a sensitivity of 92%, Specificity of 87%, and Accuracy of 93.28% were more perfect in compare with >5.

Corresponding Author:

Dr. Muna Kadhum Farhood Malla

M. B. Ch. B \ D. G. O. \ (Obstetrics and Gynecology) Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq.

Email: drmunakadhum@gmail.com

Copyright © 2025 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

One of the leading causes of female death in developing nations is cervical cancer, which accounts for one-fourth of all cases globally in India alone. As opposed to one for every 100 women of the industrialized world, it has been found that 1 in 53 Indian women suffer from a lifetime risk of having cervical cancer. The squamocolumnar junction is a representation of the transformation zone, and it is here where aberrant nuclear alterations take place, causing cellular malignancy. Since cervical cancer is virtually always associated with the human papillomavirus (HPV), it is believed that HPV is a key factor in the development of cervical cancer [1], [2].

According to histology, squamous cell carcinoma, which accounts for 70% of all cases, is the most prevalent kind of cervical cancer [3]. Even though cervical cancer endures a lengthy latent phase, it still has an elevated prevalence and incidence in resource-poor nations like India due to a poorly organized health system and a deficiency in screening programs that are capable of identifying premalignant as well as malignant lesions. Screening techniques include the Pap smear, HPV DNA test, visual examination with acetic acid (VIA), and Lugol's iodine (VILI) [4], [5].

Cytology-based screening programs are challenging to implement in low-resource settings because they depend on laboratories and expensive equipment as well as adequate assistance from trained technicians and professional employees to produce and analyze the slides. For cytology to be successful, it must be performed repeatedly throughout time. VIA has been used as an alternate screening tool to cervical cytology because to a lack of suitable cytology screening facilities and a high number of cervical cancer patients in poor nations.

It has the benefit of being reasonably priced and producing effects right away. The accuracy and dependability of VIA have, however, come under scrutiny recently. Colposcopy is still the de facto gold standard for screening, so to speak. The location, size, margins, vascularity, and extent of aberrant cervical lesions are considered during this straightforward, noninvasive OPD treatment [6], [7], [8], [9], [10], [11]. This paper was objectived to assess the outcomes of cervical cancer to see the evaluation of the effectiveness of colposcopy in Iraqi women.

2. METHODOLOGY

This paper was presented as a cross-sectional study to assess the outcomes of cervical cancer to see the evaluation of the effectiveness of colposcopy in Iraqi women where to include from 25th August 2021 to 16th July 2022, into 50 patients for ages between 25-50 years. This paper was analysed and examine of clinical features for cervical cancer by the SPSS program. This paper was contributed into Distributions of clinical features of cervical cancer patients based on ages within 25 to 50 years, where can be seen in [Table 1](#), BMI into <27.55 and >27.55, which it was be found in [Table 2](#), economic level have a Lower class where

can be cleared in Table 3, which are Middle class, and Upper class, parity which includes $\geq P4$, P1, P2, and P3, it can be shown in Table 4.

In the progress of outcomes, this paper was progressing into findings of cervical biopsy, which include Bleeding PV, Intermenstrual bleeding, Postcoital bleeding, postmenstrual bleeding, Lower abdominal pain, and Vaginal discharge, where these outcomes were got on Figure 1. To further of outcomes, this paper was highlighted into Findings of cervical biopsy based on lesion size that divide with <5 mm, >15 mm, and 5–15 mm where these findings were got in Table 5. Moreover, this paper was progressed to analyse of swede scores into cervical biopsy patients that determine <5 and $\geq$ five, which can be seen in Table 6. In addition, this study was determined with Distributions of Swede score levels in correlation with cervical biopsy patients, which include < 5 , ≥ 5 , < 7 , and $\geq$ seven. This outcome was found in Table 7. Finally, this study was accessed to outcome with Swede scores analysis based on sensitivity, specificity, and accuracy, where this finding was shown in Figure 2.

3. RESULTS AND DISCUSSION

Table 1. Distributions of Clinical Features of Cervical Cancer Patients Based on Age.

N	V	50
	Mi	0
	M	38.0000
	Std	1.03016
	Me	38.0000
	Mo	26.00 ^a
	SD	7.28431
	Var	53.061
	Ra	24.00
	Min	26.00
	Max	50.00
	S	1900.00

Table 2. Distributions of Clinical Features of Cervical Cancer Patients Based on BMI.

		Freq	P (%)	VP (%)	CP (%)
V	<27.55	15	30.0	30.0	30.0
	>27.55	35	70.0	70.0	100.0
	T	50	100.0	100.0	

Table 3. Distributions of Clinical Features of Cervical Cancer Patients Based on An Economic Level.

		Freq	P (%)	VP (%)	CP (%)
V	Lower class	16	32.0	32.0	32.0
	Middle class	26	52.0	52.0	84.0
	Upper class	8	16.0	16.0	100.0
	T	50	100.0	100.0	

Table 4. Distributions of Clinical Features of Cervical Cancer Patients Based on Parity.

		Freq	P (%)	VP (%)	CP (%)
V	$\geq P4$	11	22.0	22.0	22.0
	P1	8	16.0	16.0	38.0
	P2	12	24.0	24.0	62.0
	P3	19	38.0	38.0	100.0

	T	50	100.0	100.0	
--	---	----	-------	-------	--




Figure 1. Findings of a Cervical Biopsy.

Table 5. Findings of Cervical Biopsy Outcomes Based on Lesion Size.

		Freq	P (%)	VP (%)	CP (%)
V	<5 mm	7	14.0	14.0	14.0
	>15 mm	17	34.0	34.0	48.0
	5-15 m	26	52.0	52.0	100.0
	T	50	100.0	100.0	

Table 6. Assessments of Swede Score into Cervical Biopsy Patients.

Swede Score	Cervical Biopsy Patients, F	Cervical Biopsy Patients, Per (%)
<5	37	74%
≥5	13	26%

Table 7. Distributions of Swede Score Level in Correlation with Cervical Biopsy Patients.

Swede Score Levels	Cervical Biopsy Patients, F	Percentage (%)
< 5	10	20%
≥ 5	40	80%
< 7	15	30%
≥ 7	35	70%




Figure 2. Swede Scores Analysis Based on Sensitivity, Specificity, and Accuracy.

Discussion

Despite having a lengthy precancerous stage, where cervical cancer remains one of the factors that has affect prevalent tumors among women in underdeveloped nations. This makes it a disorder that can be prevented because there is enough opportunity for screening and treatment. But because the health system lacks the resources and appropriate screening programs to find precancerous lesions, fatality rates are quite high, rendering carcinoma cervix the main cancer-related death of females. The Swede score had defined with used by Strander et al. as well as [12] to determine the severity associated with the cervical lesion. According to [13], the specificity of a Swede score with eight or higher was 90%, and no patients with a score of 5 had lesions of CIN 2 nor worse. Additionally, they discovered that 70% with large-sized lesions with two points had histological diagnoses of CIN 2 or worse. Eight participants with CIN 2 or greater had more than a two score for the size of the lesion in this investigation. According to research [14] a Swede score for eight or above resulted in a specificity for 95% of CIN 2 or worse and a sensitivity of 38%; decreasing the cutoff to 6 increased sensitivity into 65% but at the expense of specificity (82%). Patients with lower scores exhibited larger negative predictive values (NPPs), with a 90% NPP for scores of three or fewer. The specificity in Swede score at a score for seven or above was 87% in the current investigation, while the sensitivity for high-grade lesions was 92%. The threshold was lowered to 5, which boosted the sensitivity reached 72.42% and the specificity to 82%. Using a threshold of 5 for the Swedish score, the accuracy of the value of positive prediction is 75.48%. The accuracy considerably increased to 93.28% if the cutoff for the Swede score was set at 7. The results were similar to those that Bowring et al., as well as Strander et al., had previously reported.

4. CONCLUSION

This study shows the Swede score was good as screening modalities as performance assessed of patients who have lower abdominal pain and discharge per vaginam as well as dyspareunia, abnormal uterine bleeding which colposcopy conducted into pointing of suspicious cervix with the unhealthy case. The detection of cervical cancer was represented the greatest strategy for prevention. Because it is the most prevalent cancer that may be prevented due to its lengthy preinvasive stage (CIN), it is important to make an early diagnosis for CIN by multiple screening modalities in women. In comparison between the Swede score >5 and >7, our study noticed Swede score have a sensitivity of 92%, Specificity of 87%, and Accuracy of 93.28% were more perfect in compare with >5.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Authors Contributions Statement

Name of Authors	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Dr. Muna Kadhum Farhood Malla	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Dr. Massar Raheem Abbas	✓		✓	✓				✓				✓		✓
Dr. Zena Hataf Salih	✓			✓	✓	✓			✓		✓		✓	✓

C : Conceptualization

M : Methodology

I : Investigation

R : Resources

Vi : Visualization

Su : Supervision

So : Software	D : Data Curation	P : Project
Va : Validation	O : Writing - Original Draft	administration
Fo : Formal analysis	E : Writing - Review & Editing	Fu : Funding acquisition

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study, and their voluntary consent was obtained prior to data collection.

Ethical Approval

The study was conducted in compliance with the ethical principles outlined in the Declaration of Helsinki and approved by the relevant institutional authorities.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.




REFERENCES

- [1] U. Karya, A. Zehra, and A. Rani, 'Evaluation of Swede score and Reid score to improve the predictive value of colposcopy and its correlation with histology', *Int J Reprod Contracept Obstet Gynecol*, vol. 9, no. 5, pp. 2059-2067, 2020. doi.org/10.18203/2320-1770.ijrcog20201805
- [2] J. Jeronimo and M. Schiffman, 'Colposcopy at a crossroads', *Am J Obstet Gynecol*, vol. 195, no. 2, pp. 349-353, 2006. doi.org/10.1016/j.ajog.2006.01.091
- [3] S. Aswathy, M. A. Quereshi, and B. Kurian, 'Cervical cancer screening: Current knowledge & practice among women in a rural population of Kerala', *India. Indian J Med Res*, vol. 136, no. 2, pp. 205-210, 2012.
- [4] Comprehensive cervical cancer control: A guide to essential practice. World Health Organization 2006. ISBN 978 92 4 154895 3.
- [5] A. Nessa, M. A. Hussain, and J. N. Rahman, 'Screening for cervical neoplasia in Bangladesh using visual inspection with acetic acid', *Int J Gynecol Obstet*, vol. 111, no. 2, pp. 115-118, 2010. doi.org/10.1016/j.ijgo.2010.06.004
- [6] K. O. Ajenifuja, J. C. Gage, and A. C. Adepiti, 'A population-based study of visual inspection with acetic acid (VIA) for cervical screening in rural Nigeria', *Int J Gynecol Cancer*, vol. 23, no. 3, pp. 507-512, 2013. doi.org/10.1097/IGC.0b013e318280f395
- [7] A. J. Cagle, S. Y. Hu, and J. W. Sellors, 'Use of an expanded gold standard to estimate the accuracy of colposcopy and visual inspection with acetic acid', *Int J Cancer*, vol. 126, no. 1, pp. 156-161, 2010. doi.org/10.1002/ijc.24719
- [8] R. Reid and P. Scalzi, 'Genital warts and cervical cancer. VII. An improved colposcopic index for differentiating benign papillomavirus infections from high-grade cervical intraepithelial neoplasia', *Am J Obstet Gynecol*, vol. 153, no. 6, pp. 611-618, 1985.
- [9] B. Strander, A. Ellström-Andersson, and S. Franzén, 'The performance of a new scoring system for colposcopy in detecting high-grade dysplasia in the uterine cervix', *Acta Obstet Gynecol Scand*, vol. 84, no. 10, pp. 1013-1017, 2005. doi.org/10.1111/j.0001-6349.2005.00895.x
- [10] R. D. Chaudhary, S. A. Inamdar, and C. Hariharan, 'Correlation of diagnostic efficacy of unhealthy cervix by cytology, colposcopy and histopathology in women of rural areas', *Int J Reprod Contracept Obstet Gynecol*, vol. 3, no. 1, pp. 213-218, 2014. doi.org/10.5455/2320-1770.ijrcog20140343

- [11] S. Kushwah and B. Kushwah, 'Correlation of two colposcopic indices for predicting premalignant lesions of the cervix', J Midlife Health, vol. 8, no. 3, pp. 118-123, 2017. doi.org/10.4103/jmh.JMH.22.17
- [12] N. Rodpenpear and K. Pataradool, 'The efficacy of modified Swede Colposcopic Index in the prediction of high-grade lesions and cancer of the cervix', J Gynecol Oncol, vol. 30, no. 5, 2019. doi.org/10.3802/jgo.2019.30.e78
- [13] P. Kushtagi and P. Fernandez, 'Significance of persistent inflammatory cervical smears in sexually active women of reproductive age', J Obstet Gynaecol India, vol. 52, no. 1, pp. 124-126, 2002.
- [14] J. Bowring, B. Strander, and M. Young, 'The Swede score: Evaluation of a scoring system designed to improve the predictive value of colposcopy', J Low Genit Tract Dis, vol. 14, no. 4, pp. 301-305, 2010. doi.org/10.1097/LGT.0b013e3181d77756

How to Cite: Dr. Muna Kadhum Farhood Malla, Dr. Massar Raheem Abbas, Dr. Zena Hataf Salih (2023). Cervical cancer to see the e of the effectiveness of colposcopy in iraqi women. Journal of Prevention, Diagnosis and Management of Human Diseases (JPDMHD) 3(1), 101-107. <https://doi.org/10.55529/jpdmhd.31.101.107>

BIOGRAPHIES OF AUTHORS

	Dr. Muna Kadhum Farhood Malla (M.B.Ch.B, D.G.O.) is a specialist in Obstetrics and Gynecology at the Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq. She has extensive experience in women's health, with a particular research interest in cervical cancer screening and diagnostic modalities such as colposcopy. She served as the principal investigator and corresponding author for this study. Email: drmunakadhum@gmail.com
	Dr. Massar Raheem Abbas (M.B.Ch.B, D.G.O.) is a specialist in Obstetrics and Gynecology at the Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq. Her clinical and academic interests include gynecological oncology, colposcopy, and women's reproductive health. She has contributed to several research projects focusing on the evaluation and early detection of cervical cancer in Iraqi women. Email: massaralhusseiny6@gmail.com
	Dr. Zena Hataf Salih (M.B.Ch.B, D.G.O.) is a specialist in Obstetrics and Gynecology at the Iraqi Ministry of Health, Najaf Health Department, Al-Furat Al-Awsat Teaching Hospital, Najaf, Iraq. Her professional focus is on women's reproductive health, preventive gynecology, and early cancer detection methods. She has collaborated on multiple clinical studies addressing cervical cancer, screening practices, and the effectiveness of diagnostic tools. Email: zenaalfatlawi254@gmail.com