

Research Paper



Quality of life measures for iraqi children's patients with covid-19

Dr. Ruaa Salah Mahdi^{1*}, Dr. Hasan Salih Qadoori², Dr. Hanan Nabeel Khalaf³

¹M.B.Ch.B., F.I.C.M.S. \ (Pediatric) Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth, Baghdad, Iraq.

²M.B.Ch.B., C.A.B.P. \ (Pediatric) Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth, Baghdad, Iraq.

³M.B.Ch.B., C.A.B.P. \ (Pediatric) Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth, Baghdad, Iraq.

Article Info

Article History:

Received: 28 January 2023

Revised: 11 April 2023

Accepted: 18 April 2023

Published: 04 June 2023

Keywords:


COVID-19

Quality of Life Measure

Physical Activity

Psychological Effect

Pediatric Patients



ABSTRACT

Background: The COVID-19 pandemic has had a major impact on public health in particular and on the survival rate in general.

Objective: This study aimed to assess children's quality of life during COVID-19. **Patients and methods:** A prospective study was conducted on paediatric patients infected with Covid-19, aged between 2 and 15 years, in different hospitals in Iraq, from July 17, 2022, to March 15, 2023. The study assessed the patients' general health by comparing their quality of life before and during COVID-19 using the ED-EQ scale. It also evaluated the survival rate of paediatric patients during Covid-19, using the Kaplan-Meier survival scale. **Results:** The results showed data for pediatric patients infected with COVID-19, indicating that patients in the age group (12-15) years recorded the highest infection rate, with 50 cases, with the percentage of males reaching 64 cases and females 38 cases. The most common symptoms were accompanied by fever in 21 patients and diarrhea in 19 patients. The clinical outcomes recorded that the dependent parameters in patients' health decreased during COVID-19 compared to before COVID-19, and they were general health factor (65.35 ± 7.8), physical factor (67.82 ± 9.2), and school interaction (54.20 ± 8.9). This is the most common of these findings. Moreover, these results recorded the 70-day survival rate of pediatric patients during and after COVID-19, as the survival rates were significantly lower during the period between days ten and day 60 compared to post-COVID-19. **Conclusion:** Although COVID-19 appears to be less frequent in children than in adults, it has a detrimental impact on children's quality of life, posing considerable physical, psychological, as well as mental health on children.

Corresponding Author:

Dr. Ruaa Salah Mahdi

M.B.Ch.B., F.I.C.M.S. \ (Pediatric) Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth, Baghdad, Iraq.

Email: Luaybohia126@gmail.com

Copyright © 2025 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

The pandemic due to the SARS-CoV-2 virus is having a great impact on the whole planet and, especially in our country. It is a challenge for our health system, with an important health, social, and economic impact due to the consequences for our way of life, which has the fight against disease and its morbidity and mortality [1]. The pandemic generated by COVID-19 has become one of the greatest challenges that the world's population has had to face during the present century. The pandemic due to the SARS-CoV-2 virus has caused drastic changes in our daily life, in addition to important health and socio-economic consequences [2], [3], [4]. As for the transmission mechanisms, it seems that there are viral factors that can favor infection (due to increased transmissibility, evasion of previous immunity, lower detection by diagnostic tests, and clinical severity), as demonstrated by the various SARS-CoV2 mutations suffered so far. [5], [6] this fact is evident with the new Omicron variant that, due to the change in capsular proteins, improves adherence to ACE2 receptors and, along with other possible factors, increases transmissibility [7].

The pandemic generated by COVID-19 has become one of the greatest challenges that the world's population has had to face during the present century [8]. The pandemic due to the SARS-CoV-2 virus has caused drastic changes in our daily life, in addition to important health and socio-economic consequences [9]. This fact is evident with the new Omicron variant that, due to the change in capsular proteins, improves adherence to ACE2 receptors and, along with other possible factors, increases transmissibility [10]. The new coronavirus, called SARS-CoV-2, causes the COVID-19 disease. Its severity is characterized by a massive inflammatory response resulting in multi-organ dysfunction [11]. Acute respiratory distress syndrome is a complication that needs to be treated in the intensive care unit and requires a multidisciplinary team [12]. There is currently no specific nutritional treatment for pediatric patients infected with this virus; however, nutritional therapy is aimed at the symptoms that appear. Patients who are not intubated or not hospitalized may have fever and diarrhea [13].

It is important that pediatric patients with COVID-19 maintain an adequate nutritional status that prevents them from becoming malnourished, overweight, or obese [14]. It is necessary to promote proper nutrition and avoid deficiencies of vitamins and inorganic nutrients [15]. In hospitalized patients, the previous nutritional status and the one they should have during their stay in intensive care will have an impact on the prognosis and requirement of ventilatory support and, therefore, on long-term morbidity and mortality [16]. There is a huge amount of research on COVID-19, initially carried out urgently and under exceptional conditions, which led to the publication of studies with important limitations. The information has acquired surprising dimensions for its volume, immediacy, and, progressively, for its quality and collaborative character [17], [18].

The pediatric population has been affected as much as other age groups. Most infections have been mild or moderate or severe symptoms with a small number of patients with potentially severe clinical forms [19]. In these circumstances, the Association of Primary Care Pediatrics considered it necessary to prepare a document on the available evidence of COVID-19 in the pediatric population (0-18 years), addressing all aspects of the disease: epidemiology, clinical, diagnosis, treatment, prevention, and vaccines [20].

2. METHODOLOGY

We conducted a prospective study of pediatric patients infected with COVID-19 in different hospitals in Iraq to evaluate the health quality of life and survival rate during follow-up from July 17th, 2022,

to March 15th, 2023. Comprehensive and relevant reports were collected for 102 patients infected with COVID-19 by conducting a questionnaire linked to the results of clinical examinations and COVID-19 PCR tests, which included all patients in the hospital. This study determined the clinical data parameters of Covid-19 patients who ranged between (2 - 15) years in terms of age, gender, symptoms, educational level, income level, and parent's marital status. Also, we conducted a comprehensive analysis of pediatric patients, which included examinations in terms of respiratory rate, systolic and diastolic blood pressure, oxygen saturation, and intubation. The hospitals also recorded data on admission to the pediatric intensive care unit and length of stay in the hospital. This study showed the outcomes of COVID-19 for pediatric patients, including infection, severity, and duration of COVID-19 disease in children. We included the laboratory results of the patients through taking a blood sample and determining them as WBC, lymphocytopenia, CRP, CPK, UARBC, and UAWBC, where all criteria and laboratory tests were classified on the basis of classification determined by the patient's data. Moreover, we evaluated the daily routine of pediatric patients to show the differences and variances in general health before and during COVID-19. This study was also enhanced by assessing the quality of life of patients based on the criteria of general health, emotional aspect, physical factor, social aspect, and social and school activities. Also, this study contributed to conducting a prospective evaluation of the use of the ED-EQ scale, as it was adopted to predict and determine the estimated long-term adverse health conditions among sick children infected with COVID-19. Also, this study identified risk factors affecting pediatric patients infected with Covid-19. This study was also conducted by determining the probability of pediatric patients surviving after and during COVID-19 through the Kaplan-Meier survival scale.

3. RESULTS AND DISCUSSION

Table 1. Demographic Characteristics of Children with Covid-19 in this Study

Characteristics	Number of patients [102]	Percentage [%]
Age [years]		
2- 6	20	19.61%
7-11	32	31.37%
12-15	50	49.02%
Gender		
Males	64	62.75%
Females	38	37.25%
Symptoms		
Fever	21	20.59%
Cough	17	16.67%
Diarrhea	19	18.63%
Breathing difficulties	13	12.75%
Headache	10	9.80%
Abdominal pain	16	15.69%
Stuffy or runny nose	6	5.88%
Nausea	21	20.59%
Parents education		
High school	30	29.41%
Undergraduate	24	23.53%
Postgraduate	48	47.06%
Income, \$		
300-500	33	32.35%
501-700	17	16.67%
> 700	52	50.98%
Marital status		

Married	66	64.71%
Divorced	23	22.55%
Widowed	13	12.75%
Respiratory rate		
< 25	20	19.61%
≥ 25	82	80.39%
Systolic/diastolic blood pressure		
≤ 128/88	14	13.73%
> 128/88	88	86.27%
PICU admission		
Yes	30	29.41%
No	72	70.59%
O2 saturation		
≥ 95%	67	65.69%
< 95%	35	34.31%
Intubation		
Yes	6	5.88%
No	96	94.12%
Length of hospital stay (day)		
< 6	76	74.51%
7 – 13	22	21.57%
> 14	4	3.92%

Table 2. Determine COVID-19 Data Associated With Children

Parameters	Number of patients [102]	Percentage [%]
Covid-19 infection		
A family member was infected	89	87.25%
A relative died of COVID-19	13	12.75%
Severity of covid-19		
Mild	15	14.71%
Moderate	50	49.02%
Sever	37	36.27%
Duration of covid-19, Days		
5 - 6 Days	27	26.47%
7- 12 Days	51	50.00%
13 – 17 Days	24	23.53%

Table 3. Laboratory Outcomes

Variables	Number of patients [102]	Percentage [%]
WBC ($\times 10^3 / \mu\text{L}$)		
< 5000	65	63.73%
5000 – 15000	21	20.59%
> 15000	16	15.69%
Lymphopenia		
Yes	28	27.45%
No	76	74.51%
CRP		
0	37	36.27%
+1	30	29.41%
+2	22	21.57%

+3	13	12.75%
CPK (U/L)		
≤ 200	88	86.27%
> 200	14	13.73%
UARBC		
≤4	92	90.20%
> 4	10	9.80%
UAWBC		
Normal	86	84.31%
Abnormal	16	15.69%

Table 4. Daily Active Routine of Children Before and During the Pandemic Covid-19

Variables	Before Covid-19	During Covid-19
Screen time [hours]		
< 3 hr	60 [58.82%]	37 [36.27%]
> 3hr	42 [41.18%]	65 [63.73%]
Physical activity		
No	21 [20.59%]	77 [75.49%]
Yes	81 [79.41%]	24 [24.51%]
Seep		
Poor	17 [16.67%]	35 [34.51%]
Good	85 [83.33%]	67 [65.69%]
Eating habits		
Worse	17 [16.67%]	30 [29.41%]
Good	40 [39.22%]	32 [31.37%]
Excellent	45 [44.12%]	40 [39.22%]

Table 5. Assessment of quality-life associated with health outcomes of paediatric patients before and during covid-19

Items	Before Covid-19	During Covid-19
General Health	88 ± 6.5	65.35 ± 7.8
Physical factor	92.55 ± 3.7	67.82 ± 9.2
Emotional factor	86.46 ± 6.8	67.48 ± 8.52
School Activity	90.60 ± 5.7	54.20 ± 8.9
Social factor	91.13 ± 2.4	70.69 ± 7.2


Table 6. Assessment of Estimated Long-Term Adverse Health Outcomes in Paediatric Patients with Covid-19

Parameters	Quality-life scores
Multisystem Inflammatory Syndrome (MIS), (mean ± SD)	
Pneumonia	55 ± 12.45
Shortness of breath	67.36 ± 12.54
Cough	57.86 ± 14.68
Long-lasting symptoms (mean ± SD)	
Fatigue	60.65 ± 9.5
Cognitive difficulties	70.74 ± 4.4
Headache	58.92 ± 10.4
Psychological effect (mean ± SD)	
Anxiety	77.5 ± 11.47
Post-traumatic stress	

Changes in behavior	76.54 ± 12.4
---------------------	--------------

Table 7. Multivariate Logistic Regression for Paediatric Patients with Covid-19

Risk factors	OR	(CI 95%)
Fever	2.01	0.8 - 4.16
Nausea	1.6	0.6 - 3.7
A family member was infected	3.55	1.41 - 6.85
Severity of covid-19	2.6	0.6 - 3.06
Multisystem Inflammatory Syndrome	4.6	1.44 - 7.89
Long-lasting symptoms	2.86	1.22 - 5.7
Post-traumatic stress	2.8	1.48 - 8.96

**Figure 1.** Survival-Life Rate of Pediatric Patients after and During Covid-19.

Discussion

The results recorded data on paediatric patients infected with Covid 19, showing that patients in the age group (12-15) years had the highest infection rate with 50 cases, with the percentage of male's 64 cases and females 38 cases, as the most common symptoms were accompanied by fever in 21 patients and diarrhoea in 19 patients. For the primary diagnoses, the initial examinations showed respiratory rate ≥ 25 in 82 patients, systolic/diastolic blood pressure $> 128/88$ in 88 patients, 30 children were admitted to the paediatric intensive care unit, and oxygen saturation $\geq 95\%$ in 67 patients and $< 95\%$ in 35 patients. The length of hospital stay was less than six days for 76 patients. Based on the PCR test for Covid-19 patients, one family member was infected with Covid-19. There were 89 patients, as the level of COVID-19 was found to be mild in 15 cases, moderate in 50 cases, and severe in 37 cases. Laboratory tests showed WBC < 5000 in 65 patients, lymphocytopenia in 28 patients, UARBC ≤ 4 in 92 patients, normal UAWBC in 86 patients, and above normal in 16 patients. Regarding the quality of life of the patients before and during COVID-19, the clinical results recorded that the standards adopted in the patients' health decreased during Covid compared to before COVID-19, and they were the public health factor (65.35 ± 7.8), the physical factor (67.82 ± 9.2) and the interaction in school (54.20 ± 8.9) were the most common of these results. In addition, future results of paediatric patients showed that these factors are the most common aetiological factors affecting children after injury, such as pneumonia (55 ± 12.45), cough (57.86 ± 14.68), fatigue (60.65 ± 9.5), and post-traumatic stress (62 ± 4.80). In addition, these results recorded the survival rate of paediatric patients during and after Covid-19 within 70 days, as survival rates were extremely low between the tenth and the sixtieth day compared to after Covid-19. Previous studies have shown that Covid-19 poses a major

risk to the quality of children's public and mental health. These studies confirmed that COVID-19 causes a sharp decline in the level of physical activity, which impairs the respiratory rate of children [21]. Another study indicated that COVID-19 negatively affects psychological factors, as children suffer from severe anxiety and stress, which results in a disruption in their daily routine, which poses a great threat to mental health [22]. Moreover, an American study confirmed the poor health services systems and the lack of availability of health care may make it difficult to maintain the survival rate [23].

4. CONCLUSION

Although COVID-19 appears less common than in adults, COVID-19 negatively affects children's quality of life, which poses a significant physical, psychological, and mental health burden on children. This study indicated that poor health care, diet, and low family income exacerbate the risk to children's general health, which does not allow them to cope better. The study results suggest that a decrease in oxygen levels and the presence of lymphopenia are associated with reduced survival rates. Additionally, logistic regression analysis revealed an increased hospital stay rate is a risk factor for Covid-19 complications. Poor efficiency in healthcare systems causes a decrease in patients' quality of life, which is attributed to a decrease in the survival rate during the COVID-19 pandemic. The predictive results in this study showed that Covid-19 is a risk factor for patients' long-term health in terms of physical and psychological aspects and long-term symptoms.

Acknowledgments

The authors have no specific acknowledgments to make for this research.

Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Author's Contributions Statement

Name of Authors	C	M	So	Va	Fo	I	R	D	O	E	Vi	Su	P	Fu
Dr. Ruaa Salah Mahdi	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Dr. Hasan Salih Qadoori	✓		✓	✓				✓				✓		✓
Dr. Hanan Nabeel Khalaf	✓			✓	✓	✓			✓		✓		✓	✓

C : Conceptualization

M : Methodology

So : Software

Va : Validation

Fo : Formal analysis

I : Investigation

R : Resources

D : Data Curation

O : Writing - Original Draft

E : Writing - Review & Editing

Vi : Visualization

Su : Supervision

P : Project

administration

Fu : Funding acquisition

Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

Informed Consent

All participants were informed about the purpose of the study, and their voluntary consent was obtained prior to data collection.

Ethical Approval

The study was conducted in compliance with the ethical principles outlined in the Declaration of Helsinki and approved by the relevant institutional authorities.

Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.


REFERENCES

- [1] R. Viner et al., 'Impacts of school closures on the physical and mental health of children and young people: A systematic review', *JAMA Pediatr*, vol. 176, pp. 400-409, 2022. doi.org/10.1001/jamapediatrics.2021.5840
- [2] U. Ravens-Sieberer et al., 'Quality of life and mental health in children and adolescents during the first year of the COVID-19 pandemic: Results of a two-wave nationwide population-based study', *Eur. Child Adolesc. Psychiatry*, vol. 32, pp. 575-588, 2021. doi.org/10.1007/s00787-021-01889-1
- [3] U. Ravens-Sieberer, A. Kaman, M. Erhart, J. Devine, R. Schlack, and C. Otto, 'Impact of the COVID-19 pandemic on quality of life and mental health in children and adolescents in Germany', *Eur. Child Adolesc. Psychiatry*, vol. 31, pp. 879-889, 2022. doi.org/10.1007/s00787-021-01726-5
- [4] D. Adibelli and A. Sümen, 'The effect of the coronavirus (COVID-19) pandemic on health-related quality of life in children', *Child. Youth Serv. Rev.*, vol. 119, 2020. doi.org/10.1016/j.chilcyouth.2020.105595
- [5] A. K. Ø. Geirdal et al., 'The significance of demographic variables on psychosocial health from the early stage and nine months after the COVID-19 pandemic outbreak. A Cross-National Study', *Int. J. Environ. Res. Public Health*, vol. 18, 2021. doi.org/10.3390/ijerph18084345
- [6] A. L. Shamblaw, R. L. Rumas, and M. W. Best, 'Coping during the COVID-19 pandemic: Relations with mental health and quality of life', *Can. Psychol. Psychol. Can.*, vol. 62, 2021. doi.org/10.1037/cap0000263
- [7] N. Suryavanshi et al., 'Mental health and quality of life among healthcare professionals during the COVID-19 pandemic in India', *Brain Behav.*, vol. 10, 2020. doi.org/10.1002/brb3.1837
- [8] F. Reiss, A. K. Meyrose, C. Otto, T. Lampert, F. Klasen, and U. Ravens-Sieberer, 'Socioeconomic status, stressful life situations and mental health problems in children and adolescents: Results of the German BELLA cohort-study', *PLoS ONE*, vol. 14, no. 3, 2019. doi.org/10.1371/journal.pone.0213700
- [9] E. Mahase, 'Covid-19: What New Variants Are Emerging and How Are They Being Investigated?', *British Medical Journal Publishing Group*, 2021. doi.org/10.1136/bmj.n158
- [10] T. Kirby, 'New variant of SARS-CoV-2 in UK causes surge of COVID-19', *The Lancet Respiratory Medicine*, vol. 9, no. 2, pp. e20-e21, 2021. [doi.org/10.1016/S2213-2600\(21\)00005-9](https://doi.org/10.1016/S2213-2600(21)00005-9)
- [11] D. Buonsenso, D. Munblit, and C. D. Rose, 'Preliminary evidence on long COVID in children', *Acta Paediatrica*, vol. 110, no. 7, pp. 2208-2211, 1992. doi.org/10.1111/apa.15870
- [12] D. Buonsenso, D. D. Giuda, and L. Sigfrid, 'Evidence of lung perfusion defects and ongoing inflammation in an adolescent with post-acute sequelae of SARS-CoV-2 infection', *The Lancet Child & Adolescent Health*, vol. 5, no. 9, pp. 677-680, 2021. [doi.org/10.1016/S2352-4642\(21\)00196-6](https://doi.org/10.1016/S2352-4642(21)00196-6)
- [13] E. Von Elm, D. G. Altman, M. Egger, S. J. Pocock, P. C. Gøtzsche, and J. P. Vandenbroucke, 'The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies', *Preventive Medicine*, vol. 45, no. 4, pp. 247-251, 2007. doi.org/10.1016/j.ypmed.2007.08.012
- [14] G. A. Saeed, W. Gaba, and A. Shah, 'Correlation between chest CT severity scores and the clinical parameters of adult patients with COVID-19 pneumonia', *Radiology research and practice*, vol. 2021. doi.org/10.1101/2020.10.15.20213058
- [15] World Health Organization, *Clinical Management of COVID-19: Interim Guidance*, World Health Organization. Geneva, Switzerland, 2020.
- [16] S. Madani, S. Shahin, M. Yoosefi, et al., "Red flags of poor prognosis in pediatric cases of COVID-19: the first 6610 hospitalized children in Iran," *BMC Pediatrics*, vol. 21, no. 1, pp. 563-569, 2021. doi.org/10.1186/s12887-021-03030-2

- [17] J. F. Ludvigsson, 'Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults', *Acta Paediatrica*, vol. 109, no. 6, pp. 1088-1095, 2020. doi.org/10.1111/apa.15270
- [18] E. A. Oliveira, E. A. Colosimo, and A. C. Simões E Silva, 'Clinical characteristics and risk factors for death among hospitalised children and adolescents with COVID-19 in Brazil: an analysis of a nationwide database', *The Lancet Child and Adolescent Health*, vol. 5, no. 8, pp. 559-568, 2021. [doi.org/10.1016/S2352-4642\(21\)00134-6](https://doi.org/10.1016/S2352-4642(21)00134-6)
- [19] Y. Dong, X. Mo, and Y. Hu, 'Epidemiology of COVID-19 among children in China', *Pediatrics*, vol. 145, no. 6, 2020. doi.org/10.1542/peds.2020-0702
- [20] S. Yeasmin et al., 'Impact of COVID-19 pandemic on the mental health of children in Bangladesh: a cross-sectional study', *Child Youth Ser Rev*, vol. 117, 2020. doi.org/10.1016/j.chilgyouth.2020.105277
- [21] K. Saurabh and S. Ranjan, 'Compliance and psychological impact of quarantine in children and adolescents due to COVID-19 pandemic', *Indian J Pediatr*, vol. 87, no. 7, pp. 532-536, 2020. doi.org/10.1007/s12098-020-03347-3
- [22] M. A. Garcia De Avila et al., 'Children's anxiety and factors related to the COVID-19 pandemic: an exploratory study using the children's anxiety questionnaire and the numerical rating scale', *Int J Environ Res Public Health*, vol. 17, 2020. doi.org/10.3390/ijerph17165757
- [23] J. Lee, S. S. Park, T. Y. Kim, D. G. Lee, and D. W. Kim, "Lymphopenia as a biological predictor of outcomes in COVID-19 patients: a nationwide cohort study," *Cancers*, vol. 13, no. 3, p. 471, 2021. doi.org/10.3390/cancers13030471

How to Cite: Dr. Ruaa Salah Mahdi, Dr. Hasan Salih Qadoori, Dr. Hanan Nabeel Khalaf (2023). Quality of life measures for iraqi children's patients with covid-19. *Journal of Prevention, Diagnosis and Management of Human Diseases (JPDMHD)* 3(1), 108-116. <https://doi.org/10.55529/jpdmhd.31.108.116>

BIOGRAPHIES OF AUTHORS

	Dr. Ruaa Salah Mahdi is a Pediatric Specialist (M.B.Ch.B., F.I.C.M.S.) at the Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth. Her research interests focus on pediatric health, infectious diseases, and improving the quality of life in children with chronic and acute illnesses. She has actively contributed to clinical studies related to COVID-19 in children. Email: Luaybohia126@gmail.com
	Dr. Hasan Salih Qadoori holds an M.B.Ch.B. and C.A.B.P. (Pediatrics) and works at the Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth. He specializes in child health, clinical pediatrics, and infectious disease management. His work includes evaluating clinical outcomes and pediatric survival during pandemics. Email: hassanazzawi1@gmail.com
	Dr. Hanan Nabeel Khalaf is a Pediatric Consultant (M.B.Ch.B., C.A.B.P.) affiliated with the Iraqi Ministry of Health, Baghdad Al-Karkh Health Directorate, Karkh Hospital for Child Birth. Her areas of expertise include pediatric care, clinical research, and child welfare. She has participated in several studies assessing the physical and psychological impact of COVID-19 on children. Email: hanannabeel15@gmail.com