

Research Paper



## Assessment the role of kidney function and total proteins in patients with diabetic nephropathy in kirkuk city/iraq

Rokan Hazem Hamad<sup>1\*</sup>, Sahib Jumaah Abdulrahman<sup>2</sup>

<sup>1</sup>Master Student, College of Education for Pure Sciences, University of Kirkuk, Iraq.

<sup>2</sup>Professor, College of Education for Pure Sciences, University of Kirkuk, Iraq.

### Article Info

#### Article History:

Received: 04 September 2023

Revised: 16 November 2023

Accepted: 22 November 2023

Published: 08 January 2024

### ABSTRACT

The current study aims to determine the levels of creatinine, urea, glomerular filtration rate eGFR and total proteins in patients with diabetic nephropathy, the experiment was conducted for the period from the beginning of July 2023 until the end of October 2023 Blood samples were collected from patients visiting Kirkuk General Hospital and specialized medical clinics in the city of Kirkuk, and included (70) male patients with diabetic nephropathy at the ages of (35-75) years and an average weight of (79) kg, as well as About (20) healthy people with the same average age and weight of patients and the study samples were distributed as follows: The first group of control group included (20) healthy males and the second group of patients group was distributed into four groups according to age groups age group (35-45) years and included 16 patients, age group (46-55) years and included 17 patients, age group (56-65) years and included 20 patients age group (66-75) years and included 17 patients. The study's findings indicated a noteworthy rise ( $P \leq 0.05$ ) in the levels of creatinine and urea and a significant decrease in the rate of glomerular filtration and total proteins in patients with diabetes compared to the control group, either by age groups, the results showed no significant differences in the concentration of creatinine and urea in patients with diabetes, while the rate of glomerular filtration and total proteins by age groups showed significant differences in patients with diabetes, as the first category showed a significant increase Compared to the rest of the categories.

#### Keywords:

Kidney Function

Egfr

Total Proteins

Diabetic Nephropathy

Serum Creatinine



#### Corresponding Author:

Rokan Hazem Hamad

Master Student, College of Education for Pure Sciences, University of Kirkuk, Iraq.

Email: [epbm22007@uokirkuk.edu.iq](mailto:epbm22007@uokirkuk.edu.iq)

Copyright © 2024 The Author(s). This is an open access article distributed under the Creative Commons Attribution License, (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. INTRODUCTION

Diabetes mellitus (DM) is a chronic condition that occurs as a result of the failure of the pancreas to produce enough insulin or when the body's insulin production is ineffective or insulin receptor malfunction [1]. DM diabetes is the most prevalent endocrine disease and is a serious health concern worldwide and complications occur that increase unexpectedly if the disease is not well regulated such as hyperglycemia, ketoacidosis, hyperosmolar coma, cardiovascular disease (CVD), end-stage renal failure, retinal damage that can cause blindness, nerve damage of various kinds, and microvascular damage that can be Impotence and slow healing all cause serious long-term complications as well as slow healing of wounds [2]. Diabetic nephropathy (DN) is one of the most common chronic microvascular complications of diabetics and the leading cause of end-stage kidney disease (chronic kidney disease (CKD), DN diabetic nephropathy is characterized by excessive filtration and synovuria in the early stages followed by a gradual decrease in kidney function [3]. Assessment of kidney function is important indicators in the lives of patients with kidney disease or diseases that affect kidney function [4]. Serum creatinine and serum urea are recognized as ideal markers to know the progression of diabetic nephropathy Diabetic nephropathy is clinically diagnosed with low GFR and possible hypertension, cardiovascular disease and mortality resulting from it.

Early detection of an imbalance in creatinine and urea levels in Diagnosis, prevention and development of diabetic kidney disease [5]. GFR is widely used as an indicator of kidney function and the normal range for men is about 130 ml per minute per 1.73 m<sup>2</sup> and for women 120 ml per minute per 1.73 m<sup>2</sup>, which depends on age, sex and body size means low values with age [6]. Total proteins are one of the most abundant compounds in blood serum or plasma and proteins are involved in many vital functions as proteins are used in the formation of enzymes, hormones and antibodies as well as It plays an important role in maintaining the balance of osmotic pressure and proteins are found in all body fluids but show a very high concentration in plasma, lymphatic fluids and some secretions, the total serum protein test evaluates the total amount of proteins in the blood, as many studies have shown high levels of globulin in the blood and a decrease in the ratio of albumin to globulin (A/G) in patients with infections and advanced malignant tumors, indicating their important role in the immune system and inflammation [7]. The current study aimed to evaluate the concentrations of kidney function creatinine and urea and to know the glomerular filtration rate and total proteins, which are important indicators in patients with diabetic nephropathy.

## 2. METHODOLOGY

### Study Design

The research took place from July 2023 inception through October 2023 conclusion at Kirkuk General Hospital and specialized medical clinics in Kirkuk City. It included (70) male patients with diabetic nephropathy at the ages of (35-75) years, and an average of weights (79) kg, as well as a number of healthy people with the same average age and weight of patients, and the study samples were distributed into two groups. The first group, the control group, included (20) healthy males, and the second group, the patient group, were distributed into four groups according to age groups: there were 16 patients in the 35-45 age group, 17 patients in the 46-55 age group, 20 patients in the 56-65 age group, and 17 patients in the 66-75 age group.

### Blood Samples

Blood samples were obtained from patients via a 5 ml venous draw and were then transferred into glass tubes that contained gel, as well as into vacuum tubes with gel and clot activator but no anticoagulant. The sample was left at room temperature for 30 minutes to coagulate, and the tubes were placed in a centrifuge for a duration of 15 minutes, rotating at a speed of 3000 rpm to acquire serum.

### Biochemical Tests

Determination of concentrations of a set of biochemical indicators of the studied groups, including estimation of creatinine and urea concentration in blood serum using ready-made analysis kit (Kit) from the French-origin Biolabo [8] and determination of glomerular filtration rate (GFR) from plasma concentration of endogenous substances such as creatinine, and was performed via eGFR calculator [9]. And the determination of the total protein concentration in serum samples using a ready-made analysis kit (Kit) from the manufacturer Biolabo, of French origin, this method is based on chromatography and absorption measurement Optical using an optical spectrometer [10].

### Statistical Analysis

Statistical analysis of the results was performed using the SPSS software program based on the T-test test, where the averages of patients and healthy people were compared at a significant level ( $P \leq 0.05$ ). The ANOVA test was used to compare between age groups and at a significant level ( $P \leq 0.05$ ), and the values of the variables were described as a standard deviation  $\pm$  Mean [11].

## 3. RESULTS AND DISCUSSION

### Creatinine Concentration in Diabetic Patients and Control Group

The findings in Table 1 reveal a notable elevation ( $P \leq 0.05$ ) in the levels of creatinine in patients with diabetes, as it reached  $(1.15 \pm 0.36)$  mg / 100 ml, compared to the control group that amounted to  $(0.50 \pm 0.20)$  mg / 100 ml, and the results of the current study agreed with the study [12] who found an increase in the average value of creatinine in the diabetic group compared to the healthy group. According to age groups the results in Table 2 show that there are no significant differences between the different age groups among patients with diabetes, as the concentration of creatinine in the first age group (35-45) amounted to  $(1.00 \pm 0.31)$  mg / 100 ml, compared to the second age group (46-55) amounted to  $(1.27 \pm 0.38)$  mg / 100 ml, in the third age group (56-65) it amounted to  $(1.13 \pm 0.31)$  mg / 100 ml, and in the fourth age group (66-75), which amounted to  $(1.19 \pm 0.42)$  mg / 100 ml.

The results of this study are consistent with the findings of [13] who indicated that the difference in age does not affect the level of creatinine in people with diabetes. High blood creatinine concentration in diabetics may be due to poor kidney filtering ability, causing the accumulation of nitrogenous waste, and decreased nephronal function causing high blood creatinine levels [14]. When the kidneys are unable to function properly, the level of creatinine in the blood rises abnormally [15]. Changes in serum creatinine concentration more reliably reflect changes in the glomerular filtration rate (GFR) instead of modifications in blood urea levels, and creatinine is formed naturally at a constant rate of creatine and its concentrations in the blood rely primarily on the glomerular filtration rate (GFR only) [16].

### Urea Concentration in Diabetic Patients and Control Group

The findings in Table 1 reveal a notable elevation ( $P \leq 0.05$ ) in the levels of urea in patients with diabetes, as it reached  $(43.82 \pm 9.87)$  mg / 100 ml, compared to the control group amounted to  $(29.45 \pm 4.27)$  mg / 100 ml, and these results agreed with the results of [17] with regard to the level of urea in the blood, which found an increase in the average concentration of blood urea in diabetics compared to the healthy group. According to the age groups in Table 2 the results indicate because there are no discernible variations between the different age groups among patients with diabetes, as the concentration of urea in

the first age group (35-45) amounted to  $(39.56 \pm 8.44)$  mg / 100 ml, compared to in the second age group (46-55) amounted to  $(46.35 \pm 9.87)$  mg / 100 ml, in the third age group (56-65) amounted to  $(43.30 \pm 10.35)$  mg / 100 ml, and in the fourth age group (66-75) amounted to  $(46.05 \pm 9.92)$  (mg / 100 ml). Because the concentration of urea in serum demonstrates the ratio between the urea generated by the liver and urea elimination in the urine by the renal system, an excess of urea in plasma may be caused by raised urea manufacturing, reduced urea elimination, or an amalgamation of the two; the highest concentrations occur in the setting of decreased the urea removal in the urine because of chronic kidney disease and the associated significant decrease in glomerular filtration rate.

GFR is a key medical indicator as it determines kidney purpose, individuals with reduced renal function, no matter the cause, have a decline in the glomerular filtration rate (GFR) and there is an association between the glomerular filtration rate (GFR) and the severity of kidney disease [18].

### Calculation of Glomerular Filtration Rate in Diabetic Patients and Control Group

The findings in Table 1 show a significant decrease ( $P \leq 0.05$ ) in the estimated glomerular filtration rate eGFR in patients with diabetes, as it reached  $(80.35 \pm 22.19)$  ml / min /  $1.73\text{ m}^2$  compared to the control group as it reached  $129.05 \pm 19.31$  (ml / min /  $1.73\text{ m}^2$ ). The current study's findings corresponded with the findings of [19], which showed a decrease in glomerular filtration rate in people with diabetes compared to healthy people, and all of these previous studies suggested that a low glomerular filtration rate is associated with diabetes. The estimated eGFR is not only an indicator of reduced renal function in diabetic nephropathy patients, but also in diabetic neuropathy, diabetic retinopathy, and other diabetes-related problems, hence evaluating eGFR in diabetic patients is crucial.

According to age groups, the results of the current study in Table 2 show significant differences in different age groups among patients with diabetes, as the glomerular filtration rate increased in the first age group (35-45) amounted to  $(96.93 \pm 22.35)$  ml / min /  $1.73\text{ m}^2$ , compared to in the second age group (46-55) amounted to  $(75.64 \pm 24.44)$  ml / min /  $1.73\text{ m}^2$  and in the third age group (56-65) amounted to  $(78.60 \pm 20.48)$  ml / min /  $1.73\text{ m}^2$  and in the fourth age group (66-75) amounted to  $(70.23 \pm 21.50)$  ml / min /  $1.73\text{ m}^2$ . The results of the current study agreed with the findings of [20] as they found a strong positive relationship between aging and an increased risk of low glomerular filtration rate among patients with diabetes and also stated that aging, synovuria, high uric acid in the blood, systolic hypertension, and HbA1c greater than 7% are the main factors associated with poor glomerular filtration rate.

The major pathogen responsible for the development of diabetic kidney disease is hyperglycemia. Multiple pathophysiological disorders, such as hypertension, variable glomerular tubular reflux, renal hypoxia, steopatoxicity, foot cell injury, inflammation, mitochondrial dysfunction, autophagy dysfunction, and increased sodium and hydrogen exchanger activity, contribute to progressive glomerular sclerosis and low glomerular filtration rate once hyperglycemia occurs. The major pathogen responsible for the development of diabetic kidney disease is hyperglycemia.

Multiple pathophysiological disorders, such as hypertension, variable glomerular tubular reflux, renal hypoxia, steopatoxicity, foot cell injury, inflammation, mitochondrial dysfunction, autophagy dysfunction, and increased sodium and hydrogen exchanger activity, contribute to progressive glomerular sclerosis and low glomerular filtration rate once hyperglycemia occurs [21]. Furthermore, diabetic kidney disease is silent until it is advanced. The estimated glomerular filtration rate eGFR is used to estimate the moderate to severe stages of chronic renal disease. According to studies, the risk of cardiovascular disease and death increases as early as stage III of CKD3, which corresponds to a glomerular filtration rate of less than 60 ml / min /  $1.73\text{ m}^2$  [22].

It is important to note that the average eGFR values in this study are within the normal range in the diabetic group, which indicates that the glomerular filtration rate eGFR is not a useful indicator early in detecting diabetic nephropathy DN, as it is detected and reduced in chronic kidney disease and renal decline late in the disease stage and this is consistent with the findings of study [23].

### Total Protein Concentration in Diabetic Patients and Control Group

The findings in [Table 1](#) show a significant decrease ( $P \leq 0.05$ ) in the total protein concentration in patients with diabetes, as it reached ( $7.72 \pm 0.49$ ) g /100 ml compared to the control group that amounted to ( $8.13 \pm 0.21$ ) g / 100 ml, the results of the study agreed with the results of the study [\[24\]](#), [\[25\]](#) where they found a decrease in the concentration of total protein in the serum for patients with diabetes, The results of the current study by age groups in [Table 2](#) show that there are significant differences between the different age groups in the concentration of total protein in patients with diabetes, as the total protein concentration in the first age group (35-45) amounted to ( $7.90 \pm 0.47$ ) g /100 ml), compared to in the second age group (46-55) amounted to ( $7.73 \pm 0.35$ ) g/100 ml and in the third age group (56-65) amounted to ( $7.47 \pm 0.52$ ) g /100 ml and in the fourth age group (66-75) amounted to ( $7.78 \pm 0.62$ ) g /100 ml, and the results of this study agreed with the findings of a study [\[26\]](#), which showed that low levels of total proteins in serum are associated with age and duration of diabetes and cardiovascular disease.

One possible cause of low total protein level associated with diabetics is the low glomerular filtration rate (GFR), which is the selective filtration of substances and minerals dissolved in the blood. Usually, these substances are filtered into the kidneys during primary blood purification, but are often reabsorbed by the renal tubules in later stages of the filtration process as a result of impaired kidney function [\[27\]](#). High blood sugar in diabetes also contributes to the formation of free radicals and these radicals cause oxidative stress and thus weaken the internal antioxidant defense system, if the amount of insulin available is insufficient, the cells respond poorly to the effects of insulin leading to continued elevation of blood sugar levels, and reduced total protein synthesis in serum [\[24\]](#).

**Table 1.** Concentration Creatinine, Urea, Glomerular Filtration Rate and Total Proteins in Diabetics and Control Group

| Variables     | Group | Patient                | Control                 |
|---------------|-------|------------------------|-------------------------|
| Creatinine    |       | $1.15 \pm 0.36$<br>a   | $0.50 \pm 0.20$<br>b    |
| Urea          |       | $43.81 \pm 9.64$<br>a  | $29.45 \pm 4.27$<br>b   |
| eGFR          |       | $80.35 \pm 22.19$<br>a | $129.05 \pm 19.31$<br>b |
| Total Protein |       | $7.72 \pm 0.49$<br>a   | $8.13 \pm 0.21$<br>b    |

\*The values in the table indicate to (Mean  $\pm$  S.D)

\*Different letters horizontally indicate significant differences at ( $P \leq 0.05$ )

**Table 2.** Concentration Creatinine, Urea, Glomerular Filtration Rate and Total Proteins by Age Groups in Diabetics

| Age Group Variables | Age ( 35 - 45 )        | Age ( 46 - 55 )        | Age ( 56 - 65 )        | Age ( 66 - 75 )        |
|---------------------|------------------------|------------------------|------------------------|------------------------|
| Creatinine          | $1.00 \pm 0.31$<br>a   | $1.27 \pm 0.38$<br>a   | $1.13 \pm 0.31$<br>a   | $1.19 \pm 0.42$<br>a   |
| Urea                | $39.56 \pm 8.44$<br>a  | $46.35 \pm 9.87$<br>a  | $43.30 \pm 10.35$<br>a | $46.05 \pm 9.92$<br>a  |
| eGFR                | $96.93 \pm 22.35$<br>a | $75.64 \pm 24.44$<br>b | $78.60 \pm 20.48$<br>b | $70.23 \pm 21.50$<br>b |
| Total Protein       | $7.90 \pm 0.47$<br>a   | $7.73 \pm 0.35$<br>b   | $7.47 \pm 0.52$<br>b   | $7.78 \pm 0.62$<br>b   |

\*The values in the table indicate to (Mean  $\pm$  S.D)

\*Different letters horizontally indicate significant differences at ( $P \leq 0.05$ ).

#### 4. CONCLUSION

From the present study, we conclude that there is a moral rise ( $P \leq 0.05$ ) in the concentration of creatinin and urea among diabetic patients compared to healthy, a decrease in the concentration of total protein and the glomerular filtration rate of diabetes patients compared to healthy. By age group, however, we conclude that there are no moral differences ( $P \leq 0.05$ ) in the concentration of creatinine and urea among diabetic patients, and that there are moral differences in the concentration of total proteins and glomerular filtration rate by age groups among diabetic patients.

#### Acknowledgments

The authors have no specific acknowledgments to make for this research.

#### Funding Information

This research received no specific grant from any funding agency in the public, commercial, or not for profit sectors.

#### Author Contributions Statement

| Name of Author           | C | M | So | Va | Fo | I | R | D | O | E | Vi | Su | P | Fu |
|--------------------------|---|---|----|----|----|---|---|---|---|---|----|----|---|----|
| Rokan Hazem Hamad        | ✓ | ✓ | ✓  | ✓  | ✓  | ✓ |   |   | ✓ | ✓ | ✓  | ✓  | ✓ |    |
| Sahib Jumaah Abdulrahman |   | ✓ |    | ✓  |    | ✓ |   | ✓ | ✓ |   |    | ✓  |   | ✓  |

C: Conceptualization

M: Methodology

So: Software

Va: Validation

Fo: Formal analysis

I: Investigation

R: Resources

D: Data Curation

O: Writing- Original Draft

E: Writing- Review & Editing

Vi: Visualization

Su: Supervision

P: Project administration

Fu: Funding acquisition

#### Conflict of Interest Statement

The authors declare that there are no conflicts of interest regarding the publication of this paper.

#### Informed Consent

All participants were informed about the purpose of the study, and their voluntary consent was obtained prior to data collection.

#### Ethical Approval

The study was conducted in compliance with the ethical principles outlined in the Declaration of Helsinki and approved by the relevant institutional authorities.

#### Data Availability

The data that support the findings of this study are available from the corresponding author upon reasonable request.

## REFERENCES

- [1] N. M. Jasem and A. S. Abdul-Razaq, 'Evalution of Biomarkers in Iraq Patients with Diabetes Mellitus Type 2', *The Egyptian Journal of Hospital Medicine*, vol. 90, no. 2, pp. 3062-3066, 2023. [doi.org/10.21608/ejhm.2023.288392](https://doi.org/10.21608/ejhm.2023.288392)
- [2] World Health Organization. *World health statistics overview 2019: monitoring health for the SDGs, sustainable development goals* (No. WHO/DAD/2019.1). World Health Organization.
- [3] M. K. Sagoo and L. Gnudi, 'Diabetic Nephropathy: An Overview', in *Diabetic Nephropathy*, New York, NY: Springer US, 2020, pp. 3-7. [doi.org/10.1007/978-1-4939-9841-8\\_1](https://doi.org/10.1007/978-1-4939-9841-8_1)
- [4] R. N. Okoro and V. T. Farate, 'The use of nephrotoxic drugs in patients with chronic kidney disease', *Int. J. Clin. Pharm.*, vol. 41, no. 3, pp. 767-775, June 2019. [doi.org/10.1007/s11096-019-00811-9](https://doi.org/10.1007/s11096-019-00811-9)
- [5] P. Lalla, S. Banu Pathan, and P. Jawade, 'Correlation of Serum Urea and Serum Creatinine in Diabetics patients and normal individuals', *Int. J. Clin. Biochem. Res.*, vol. 7, no. 1, pp. 45-48, May 2020. [doi.org/10.18231/ijcbr.2020.009](https://doi.org/10.18231/ijcbr.2020.009)
- [6] L. A. Stevens, J. Coresh, T. Greene, and A. S. Levey, 'Assessing kidney function--measured and estimated glomerular filtration rate', *N. Engl. J. Med.*, vol. 354, no. 23, pp. 2473-2483, June 2006. [doi.org/10.1056/NEJMra054415](https://doi.org/10.1056/NEJMra054415)
- [7] Y. Yoshino et al., 'A low albumin to globulin ratio with a high serum globulin level is a prognostic marker for poor survival in cervical cancer patients treated with radiation based therapy', *Int. J. Gynecol. Cancer*, vol. 29, no. 1, pp. 17-22, Jan. 2019. [doi.org/10.1136/ijgc-2018-000025](https://doi.org/10.1136/ijgc-2018-000025)
- [8] S. A. Hussein, H. A. Fadlalmola, S. M. Salama, E. G. Osman, and A. A. Mariod, 'Efficacy and Safety of Gum Arabic on Renal Failure Patients: Systematic Review and Meta-analysis', *Sudan Journal of Medical Sciences*, vol. 17, no. 4, pp. 459-475, 2022. [doi.org/10.18502/sjms.v17i4.12547](https://doi.org/10.18502/sjms.v17i4.12547)
- [9] A. S. Levey and L. A. Stevens, 'Estimating GFR using the CKD Epidemiology Collaboration (CKD-EPI) creatinine equation: more accurate GFR estimates, lower CKD prevalence estimates, and better risk predictions', *Am. J. Kidney Dis.*, vol. 55, no. 4, pp. 622-627, Apr. 2010. [doi.org/10.1053/j.ajkd.2010.02.337](https://doi.org/10.1053/j.ajkd.2010.02.337)
- [10] V. Narwal, N. Sharma, R. Sharma, Y. S. Rajput, and B. Mann, 'Applicability of protein estimation methods for assaying glycomacropeptide', *International Journal of Dairy Technology*, vol. 71, no. 2, pp. 539-543, 2018. [doi.org/10.1111/1471-0307.12452](https://doi.org/10.1111/1471-0307.12452)
- [11] A.-R. Kh, *Introduction to Statistics*, Second Edition, College of Agriculture and Forestry. 2000.
- [12] H. A. El-Attar, M. M. El-Deeb, and L. A. El-Ghlied, 'Serum Glycoprotein Chondrex (YKL-40) and High Sensitivity C-Reactive Protein (hsCRP) in Type 2 Diabetic Patients in Relation to Cardiovascular Complications', *J Nephrol Kidney Dis*, vol. 1, no. 1, 2017. [doi.org/10.36876/smjnk.1003](https://doi.org/10.36876/smjnk.1003)
- [13] N. H. Ibrahim, *Investigation of Myonectin and Irisin levels and a number of physiological criteria in people with type II diabetes in the city of Kirkuk*. 2022.
- [14] N. Yadav, S. Chandra, K. Singh, A. Yadav, and A. Sharma, 'Assessment of Prognostic markers of Diabetic Nephropathy-Serum Creatinine and Blood Urea Levels in Diabetes mellitus and healthy individuals at tertiary care hospital', *European Journal of Molecular & Clinical Medicine (EJMCM)*, vol. 7, no. 11, 2020.
- [15] A. A. Abdulrazzaq, 'Renal Impairment in Diabetes Mellitus Type2 In Iraq', *International Journal of Pharmaceutical Research*, vol. 12, no. 1, 2020. [doi.org/10.31838/ijpr/2020.12.01.329](https://doi.org/10.31838/ijpr/2020.12.01.329)
- [16] K. A. Griffin, H. Kramer, and A. K. Bidani, 'Adverse renal consequences of obesity', *Am. J. Physiol. Renal Physiol.*, vol. 294, no. 4, pp. F685-96, Apr. 2008. [doi.org/10.1152/ajprenal.00324.2007](https://doi.org/10.1152/ajprenal.00324.2007)
- [17] H. F. Hassan, *The Relation of Oxidative Stress with the Activity of Adenosine Deaminase and Its Isoenzymes in Iraqi Patients with Diabetic Kidney Disease*. Thesis. 2022. [doi.org/10.53730/ijhs.v6nS8.13342](https://doi.org/10.53730/ijhs.v6nS8.13342)
- [18] C. Higgins, 'Urea and the clinical value of measuring blood urea concentration', *Acute care testing. Org*, vol. 2016, pp. 1-6.

[19] A. Ramezankhani, F. Azizi, and F. Hadaegh, 'Association between estimated glomerular filtration rate slope and cardiovascular disease among individuals with and without diabetes: a prospective cohort study', *Cardiovasc. Diabetol.*, vol. 22, no. 1, p. 270, Oct. 2023. [doi.org/10.1186/s12933-023-02008-x](https://doi.org/10.1186/s12933-023-02008-x)

[20] N. Nata, R. Rangsin, O. Supasyndh, and B. Satirapoj, 'Impaired glomerular filtration rate in type 2 diabetes mellitus subjects: A nationwide cross-sectional study in Thailand', *J. Diabetes Res.*, vol. 2020, p. 6353949, Aug. 2020. [doi.org/10.1155/2020/6353949](https://doi.org/10.1155/2020/6353949)

[21] R. A. DeFronzo, W. B. Reeves, and A. S. Awad, 'Pathophysiology of diabetic kidney disease: impact of SGLT2 inhibitors', *Nat. Rev. Nephrol.*, vol. 17, no. 5, pp. 319-334, May 2021. [doi.org/10.1038/s41581-021-00393-8](https://doi.org/10.1038/s41581-021-00393-8)

[22] P. Sharma et al., 'Does stage-3 chronic kidney disease matter?: A systematic literature review', *British Journal of General Practice*, vol. 60, no. 575, pp. e266-e276, 2010. [doi.org/10.3399/bjgp10X502173](https://doi.org/10.3399/bjgp10X502173)

[23] S. I. Abid, NeutrophilGelatinase-associated Lipocalin (NGAL) and Asprosin in Early Detection of Nephropathy in Type2 Diabetes. Thesis. 2021.

[24] F. A. Nazki, A. Syyeda, and S. Mohammed, 'Total proteins, albumin and HbA1c in type 2 diabetes mellitus', *Medpulse Int J Biochem*, vol. 3, no. 3, pp. 40-42, 2017.

[25] M. Adnan Khalaf and I. Ghassan Zainal, 'Investigation of antioxidant markers in diabetic patients', *Arch. Razi Inst.*, vol. 76, no. 5, pp. 1453-1460, Nov. 2021.

[26] S. Riaz, M. Tariq, and S. Aslam, 'Association of serum protein levels in the diabetic patients with risk of cardiovascular disease and nephropathy in Pakistani population', *J Res Diabetes Metab*, vol. 4, no. 1, pp. 11-015, 2018.

[27] Christensson et al., 'The impact of the glomerular filtration rate on the human plasma proteome', *Proteomics Clin. Appl.*, vol. 12, no. 3, p. 1700067, May 2018. [doi.org/10.1002/prca.201700067](https://doi.org/10.1002/prca.201700067)

**How to Cite:** Rokan Hazem Hamad, Sahib Jumaah Abdulrahman. (2024). Assessment the role of kidney function and total proteins in patients with diabetic nephropathy in kirkuk city/iraq. *Journal of Prevention, Diagnosis and Management of Human Diseases (JPDMHD)*, 4(1), 13-21. <https://doi.org/10.55529/jpdmhd.41.13.21>

## BIOGRAPHIES OF AUTHORS

|   |  |
|---|--|
|  | <p><b>Rokan Hazem Hamad</b>, is a Master's student in the College of Education for Pure Sciences at the University of Kirkuk, Kirkuk, Iraq. His academic and research interests focus on clinical biochemistry, diabetes mellitus, diabetic nephropathy, and kidney function biomarkers. His current research emphasizes the evaluation of renal function parameters and biochemical indicators associated with diabetic complications. He has participated in several laboratory-based studies related to metabolic and renal disorders and aims to contribute to early diagnosis and management strategies for chronic diseases. Email: <a href="mailto:epbm22007@uokirkuk.edu.iq">epbm22007@uokirkuk.edu.iq</a></p> |
|  | <p><b>Sahib Jumaah Abdulrahman</b>, is a Professor in the College of Education for Pure Sciences at the University of Kirkuk, Kirkuk, Iraq. He has extensive teaching and research experience in clinical biochemistry and human physiology. His research interests include diabetes mellitus, renal diseases, oxidative stress, and biochemical biomarkers in chronic disorders. He has supervised numerous postgraduate theses and published several scientific papers in national and international journals. His work contributes significantly to advancing biomedical</p>  |

|  |   |
|--|---|
|  | research and academic development in Iraq. Email:<br><a href="mailto:drsahib68@uokirkuk.edu.iq">drsahib68@uokirkuk.edu.iq</a> |
|--|---|